bs-8869R

[Primary Antibody]

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

IP3 receptor Rabbit pAb

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GeneID: 3708 SWISS: Q14643

Target: IP3 receptor

Immunogen: KLH conjugated synthetic peptide derived from human IP3R 1:

101-200/2758.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: This gene encodes an intracellular receptor for inositol 1,4,5-

trisphosphate. Upon stimulation by inositol 1,4,5-trisphosphate, this receptor mediates calcium release from the endoplasmic reticulum. Mutations in this gene cause spinocerebellar ataxia type 15, a disease associated with an heterogeneous group of cerebellar disorders. Multiple transcript variants have been identified for this

gene. [provided by RefSeq, Nov 2009]

Applications: Flow-Cyt (2ug/Test)

ICC/IF (1:100-500)


Reactivity: Human (predicted: Mouse,

Rat, Rabbit, Pig, Sheep, Cow, Chicken, Dog, Horse)


Predicted 314 kDa MW.:

Subcellular Cytoplasm

VALIDATION IMAGES



HepG2 cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Antibody incubation with (IP3 receptor) polyclonal Antibody, Unconjugated (bs-8869R) 1:100, 90 minutes at 37°C; followed by a conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei.



Blank control (black line) :SH-SY5Y. Primary Antibody (green line): Rabbit Anti-IP3 receptor antibody (bs-8869R) Dilution:2ug/Test; Secondary Antibody (white blue line): Goat anti-rabbit IgG-AF488 Dilution: 0.5ug/Test. Isotype control (orange line): Normal Rabbit IgG Protocol The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 90% ice-cold methanol for 20 min at -20°C. The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

— SELECTED CITATIONS —

• [IF=2.4] Zhi-Yong An. et al. Eicosatrienoic acid enhances the quality of in vitro matured porcine oocytes by reducing PRKN-mediated ubiquitination of CISD2. THERIOGENOLOGY. 2024 Dec;230:285 IF; Pig. 39357167