
phospho-GATA3 (Ser308) Rabbit pAb

Catalog Number: bs-13294R

Target Protein: phospho-GATA3 (Ser308)

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000)

Reactivity: Human, Mouse (predicted:Rabbit, Sheep, Cow, Dog, GuineaPig)

Predicted MW: 49 kDa

Entrez Gene: 2625

Swiss Prot: P23771

Source: KLH conjugated synthesised phosphopeptide derived from human GATA3 around the phosphorylation site of Ser308: RL(pS)AA.


Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.


Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: Members of the GATA family share a conserved zinc finger DNA-binding domain and are capable of binding the WGATAR consensus sequence. GATA-1 is erythroid-specific and is responsible for the regulated transcription of erythroid genes. It is an essential component in the generation of the erythroid lineage. GATA-2 is expressed in embryonic brain and liver, HeLa and endothelial cells, as well as in erythroid cells. Studies with a modified GATA consensus sequence, AGATCTTA, have shown that GATA-2 and GATA-3 recognize this mutated consensus while GATA-1 has poor recognition of this sequence. This indicates broader regulatory capabilities of GATA-2 and GATA-3 than GATA-1. GATA-3 is highly expressed in T lymphocytes. GATA-4, GATA-5 and GATA-6 comprise a subfamily of transcription factors. Both GATA-4 and GATA-6 are found in heart, pancreas and ovary; lung and liver tissues exhibit GATA-6, but not GATA-4 expression. GATA-5 expression has been observed in differentiated heart and gut tissues and is present throughout the course of development in the heart. Although expression patterns of the various GATA transcription factors may overlap, it is not yet apparent how the GATA factors are able to discriminate in binding their appropriate target sites.

VALIDATION IMAGES



Protein: bladder(mouse) lysates at 40ug; Primary: Rabbit Anti- p-GATA3 (Ser308) (bs-13294R) at 1:300; Secondary: 800CW Conjugated Goat (polyclonal) Anti-Rabbit IgG(H+L) at 1: 10000; Predicted band size:49 kD
Observed band size:49 kD



Sample: Lane 1: Human SH-SY5Y cell lysates Lane 2: Human A431 cell lysates Primary: Anti-phospho-GATA3 (Ser308) (bs-13294R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 49 kDa Observed band size: 49 kDa