

**bs-23864R****[ Primary Antibody ]****Neuropilin 1 Rabbit pAb****Bioss**  
**ANTIBODIES**

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

**— DATASHEET —**

<b>Host:</b> Rabbit <b>Clonality:</b> Polyclonal <b>GeneID:</b> 8829 <b>Target:</b> Neuropilin 1 <b>Immunogen:</b> KLH conjugated synthetic peptide derived from human Neuropilin 1 : 101-200/923. < Extracellular > <b>Purification:</b> affinity purified by Protein A <b>Concentration:</b> 1mg/1ml <b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. <b>Background:</b> This gene encodes one of two neuropilins, which contain specific protein domains which allow them to participate in several different types of signaling pathways that control cell migration. Neuropilins contain a large N-terminal extracellular domain, made up of complement-binding, coagulation factor V/VIII, and meprin domains. These proteins also contains a short membrane-spanning domain and a small cytoplasmic domain. Neuropilins bind many ligands and various types of co-receptors; they affect cell survival, migration, and attraction. Some of the ligands and co-receptors bound by neuropilins are vascular endothelial growth factor (VEGF) and semaphorin family members. Several alternatively spliced transcript variants that encode different protein isoforms have been described for this gene. [provided by RefSeq, Oct 2011]	<b>Isotype:</b> IgG <b>SWISS:</b> O14786	<b>Applications:</b> WB (1:500-2000) <b>Reactivity:</b> Mouse, Rat (predicted: Human, Rabbit, Pig, Sheep, Cow, Zebrafish, Chicken, Dog, Horse) <b>Predicted MW.:</b> 100 kDa <b>Subcellular Location:</b> Secreted ,Cell membrane
---	---	---

**— VALIDATION IMAGES —**

Sample: Fetal brain (Mouse) Lysate at 40 ug  
Cerebrum (Mouse) Lysate at 40 ug Cerebrum  
(Rat) Lysate at 40 ug Primary: Anti-Neuropilin 1 (bs-23864R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 100 kD Observed band size: 110 kD



Sample: Liver (Mouse) Lysate at 40 ug Primary: Anti- Neuropilin 1 (bs-23864R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 100 kD Observed band size: 140 kD