

bs-12335R**[Primary Antibody]****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

HFE Rabbit pAb**— DATASHEET —**

Host: Rabbit Clonality: Polyclonal Target: HFE Immunogen: KLH conjugated synthetic peptide derived from Human HFE/Hemochromatosis: 262-348/348. Purification: affinity purified by Protein A Concentration: 1mg/ml Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. Background: The features of hemochromatosis include cirrhosis of the liver, diabetes, hypermelanotic pigmentation of the skin, and heart failure. Since hemochromatosis is a relatively easily treated disorder if diagnosed, this is a form of preventable cancer. The HFE protein, which is defective in hereditary hemo-chromatosis, normally is expressed in crypt enterocytes of the duodenum where it has a unique, predominantly intracellular localization. In placenta, the HFE protein co-localizes with and forms a stable association with the transferrin receptor (TfR), providing a link between the HFE protein and iron transport. Immunocytochemistry shows that the HFE protein and TfR both are expressed in the crypt enterocytes. Western blots show that, as is the case in human placenta, the HFE protein in crypt enterocytes is physically associated with the TfR and with β 2-microglobulin. It is proposed that HFE has two mutually exclusive activities in cells: inhibition of uptake or inhibition of release of iron and that the balance between serum transferrin saturation and serum transferrin-receptor concentrations determines which of these functions predominates. The gene which encodes HFE maps to human chromosome 6p21.3.	Isotype: IgG GenelD: 3077 Applications: Flow-Cyt (1 μ g/Test) Reactivity: Human (predicted: Mouse, Rat, Sheep, Cow, Dog, Horse) Predicted MW: 38 kDa Subcellular Location: Cell membrane
---	---

— VALIDATION IMAGES —

Blank control (blue line): HL60(fixed with 70% ethanol Overnight at 4°C). Primary Antibody (green line): Rabbit Anti-iHFE antibody (bs-12335R), Dilution: 0.2 μ g / 10⁶ cells; Isotype Control Antibody (orange line): Rabbit IgG . Secondary Antibody (white blue line): Goat anti-rabbit IgG-PE, Dilution: 1 μ g /test.

— SELECTED CITATIONS —

- **[IF=5.008]** Rychtarcikova, Zuzana, et al. "Tumor-initiating cells of breast and prostate origin show alterations in the expression of genes related to iron metabolism." Oncotarget (2016). WB ;="Human". 28031527