

PGC1 alpha + beta Rabbit pAb

Catalog Number: bs-7535R

Target Protein: PGC1 alpha + beta

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000), IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500), Flow-Cyt (1µg/Test)

Reactivity: Human, Rat (predicted: Mouse, Rabbit, Pig, Cow, Dog, Horse)

Predicted MW: 88/113 kDa

Entrez Gene: 10891

Swiss Prot: Q9UBK2

Source: KLH conjugated synthetic peptide derived from human PGC1 alpha + beta: 151-250/798.


Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.


Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: The protein encoded by this gene is a transcriptional coactivator that regulates the genes involved in energy metabolism. This protein interacts with PPARgamma, which permits the interaction of this protein with multiple transcription factors. This protein can interact with, and regulate the activities of cAMP response element binding protein (CREB) and nuclear respiratory factors (NRFs). It provides a direct link between external physiological stimuli and the regulation of mitochondrial biogenesis, and is a major factor that regulates muscle fiber type determination. This protein may be also involved in controlling blood pressure, regulating cellular cholesterol homeostasis, and the development of obesity (referenced from entrez gene).


VALIDATION IMAGES




Sample: Lane 1: Molt-4 (Human) Cell Lysate at 30 ug Lane 2: A431 (Human) Cell Lysate at 30 ug Lane 3: HepG2 (Human) Cell Lysate at 30 ug Lane 4: HeLa (Human) Cell Lysate at 30 ug Lane 5: Jurkat (Human) Cell Lysate at 30 ug Lane 6: HL60 (Human) Cell Lysate at 30 ug Primary: Anti-PGC1 alpha + beta (bs-7535R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 90/113 kDa Observed band size: 113 kDa



Tissue/cell: rat heart tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-PGC1 alpha+beta Polyclonal Antibody, Unconjugated (bs-7535R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody (SP-0023) and DAB (C-0010) staining



Paraformaldehyde-fixed, paraffin embedded (rat kidney tissue); Antigen retrieval by boiling in sodium citrate buffer (pH 6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (PGC1 alpha + beta) Polyclonal Antibody, Unconjugated (bs-7535R) at 1:400 overnight at 4°C, followed by operating according to SP Kit (Rabbit) (sp-0023) instructions and DAB staining.



Blank control (Black line): Molt4 (Black). Primary Antibody (green line): Rabbit Anti-PGC1 alpha+beta antibody (bs-7535R) Dilution: 1μg / 10⁶ cells; Isotype Control Antibody (orange line): Rabbit IgG. Secondary Antibody (white/blue line): Goat anti-rabbit IgG-AF647 Dilution: 1μg / test. Protocol: The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 90% ice-cold methanol for 20 min at room temperature. The cells were then incubated in 5% BSA to block non-specific protein-protein interactions for 30 min at room temperature. Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody was used for 40 min at room temperature. Acquisition of 20,000 events was performed.

PRODUCT SPECIFIC PUBLICATIONS

[IF=5.26] Meng, Guoliang, et al. "Hydrogen sulfide pretreatment improves mitochondrial function in myocardial hypertrophy via a SIRT3 dependent manner." *British Journal of Pharmacology* (2017). WB ; ="Mouse" . 28503736

[IF=5.279] Lei Wang, et al. Lactobacillus plantarum DP189 Reduces α-SYN Aggravation in MPTP-Induced Parkinson's Disease Mice via Regulating Oxidative Damage, Inflammation, and Gut Microbiota Disorder. *J Agr Food Chem*. 2022;70(4):1163–1173 WB ; Mouse . 35067061

[IF=4.8] Song-Xia Lin, et al. Eriodictyol regulates white adipose tissue browning and hepatic lipid metabolism in high fat diet-induced obesity mice via activating AMPK/SIRT1 pathway. *J ETHNOPHARMACOL*. 2024 Aug;:118761 WB ; Mouse . 39216775

[IF=4.315] Ma, Dufang, et al. Astragalus polysaccharide prevents heart failure-induced cachexia by alleviating excessive adipose expenditure in white and brown adipose tissue. *LIPIDS HEALTH DIS*. 2023 Dec;22(1):1-16 WB ; Rat . 36670439

[IF=3.776] Tingting Zhao, et al. Yishen Huashi granule modulated lipid metabolism in diabetic nephropathy via PI3K/AKT/mTOR signaling pathways.. *HELIYON*. 2023 Mar;9(3):e14171-e14171 WB ; Human . 36938470