

bs-1665R**[Primary Antibody]**

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

VEGFA Rabbit pAb**— DATASHEET —**

Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000)
Clonality: Polyclonal		Reactivity: Mouse, Rat
GeneID: 7422	SWISS: P15692	
Target: VEGFA		Predicted MW.: 24 kDa
Immunogen: KLH conjugated synthetic peptide derived from human VEGF: 81-132/232.		Subcellular Location: Secreted
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: Vascular endothelial growth factor (VEGF), originally known as vascular permeability factor (VPF), is a signal protein produced by cells that stimulates the formation of blood vessels. To be specific, VEGF is a sub-family of growth factors, the platelet-derived growth factor family of cystine-knot growth factors. They are important signaling proteins involved in both vasculogenesis (the de novo formation of the embryonic circulatory system) and angiogenesis (the growth of blood vessels from pre-existing vasculature).		

— SELECTED CITATIONS —

- **[IF=15.304]** Rajendra K. Singh. et al. Diabetic bone regeneration with nanoceria-tailored scaffolds by recapitulating cellular microenvironment: Activating integrin/TGF- β co-signaling of MSCs while relieving oxidative stress. BIOMATERIALS. 2022 Aug;;121732 IF ;Rat. 36031457
- **[IF=7.7]** Jingjunjiao Long. et al. Nanosilicate-reinforced GelMA-PEGDA hydrogel promotes angiogenesis for bone regeneration. INT J BIOL MACROMOL. 2024 Jun;;133202 WB ;Human. 38889828
- **[IF=5.168]** Gu et al. Fasudil attenuates soluble fms-like tyrosine kinase-1 (sFlt-1)-induced hypertension in pregnant mice through RhoA/ROCK pathway. (2017) Oncotarget. 8:104104-104112 WB,IHC ;Human, Mouse. 29262624
- **[IF=4.6]** Ke Minhui. et al. Establishment and study of a rat internal haemorrhoid model. SCI REP-UK. 2023 Dec;13(1):1-10 IHC ;Rat. 38049459
- **[IF=5.4]** Feng Qiu. et al. The mechanism of Chebulae Fructus Immaturus promote diabetic wound healing based on network pharmacology and experimental verification. J ETHNOPHARMACOL. 2024 Mar;322:117579 IHC,WB ;Mouse,Human. 38104882