[Primary Antibody]

phospho-NFKB p65 (Ser536) Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

— DATASHEET ———		400-901-9800
Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000)
Clonality: Polyclonal		IHC-P (1:50-200)
GenelD: 5970	SWISS: Q04206	IF (1:50-200)
Target: NFKB p65 (Ser536)		Reactivity: Mouse (predicted: Human, Rat, Monkey)
Immunogen: KLH conjugated Synthesised phosphopeptide derived from human NFKBp65 around the phosphorylation site of Ser536: FS(p-S)IA.		
Purification: affinity purified	l by Protein A	
Concentration: 1mg/ml		Predicted MW.: ^{61 kDa}
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Subcellular Location: ^C ytoplasm ,Nucleus
Background: NF-kappa-B is a biological proc by specific inhi B moves to the genes. NF-kapp either REL, REL is NFKB1 comp transcript varia this gene. [prov	a ubiquitous transcription factor involved in several esses. It is held in the cytoplasm in an inactive state bitors. Upon degradation of the inhibitor, NF-kappa- nucleus and activates transcription of specific ba-B is composed of NFKB1 or NFKB2 bound to A, or RELB. The most abundant form of NF-kappa-B lexed with the product of this gene, RELA. Four ints encoding different isoforms have been found for <i>i</i> ded by RefSeq, Sep 2011].	

— VALIDATION IMAGES



NIH/3T3 (M) cells were treated with or without Calyculin A (100nM) for 30 min, 25 µg total protein per lane of cell lysates (see on figure) probed with phospho-NFKB p65 polyclonal antibody, unconjugated (bs-0982R) at 1:2000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.

- SELECTED CITATIONS ------

- [IF=17.521] Yi Yan. et al. Nanomedicines Reprogram Synovial Macrophages by Scavenging Nitric Oxide and Silencing CA9 in Progressive Osteoarthritis. Advanced Science. 2023 Feb;:2207490 WB ;Mouse. 36748885
- [IF=15.304] Sitong Liu. et al. MRI-visible mesoporous polydopamine nanoparticles with enhanced antioxidant capacity for osteoarthritis therapy. BIOMATERIALS. 2023 Apr;295:122030 WB ;Mouse. 36758340
- [IF=12.4] Manikandan Santhanam. et al. Interaction of SMAC with a survivin-derived peptide alters essential cancer hallmarks: Tumor growth, inflammation, and immunosuppression. MOL THER. 2024 Apr 05 WB ;MOUSE. 38582961
- **[IF=11.7]** Ran Cheng. et al.Intratumoral antigen-presenting cell activation by a nanovesicle for the concurrent tertiary lymphoid structure de novo neogenesis.science advances.2025 Feb 21;11(8):eadr1299. ;MOUSE. 39970209

 [IF=10.684] Chen Zhang. et al. The novel hyaluronic acid granular hydrogel attenuates osteoarthritis progression by inhibiting the TLR-2/NF-κB signaling pathway through suppressing cellular senescence. BIOENG TRANSL MED. 2022
Dec;:e10475 WB ;MOUSE. 10.1002/btm2.10475