

**bs-4937R****[ Primary Antibody ]****Cystatin A Rabbit pAb**

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

**— DATASHEET —**

<b>Host:</b> Rabbit	<b>Isotype:</b> IgG	<b>Applications:</b> <b>WB</b> (1:500-2000) <b>IHC-P</b> (1:100-500) <b>IHC-F</b> (1:100-500) <b>IF</b> (1:100-500) <b>ELISA</b> (1:5000-10000)
<b>Clonality:</b> Polyclonal		<b>Reactivity:</b> (predicted: Mouse, Rat, Rabbit)
<b>Target:</b> Cystatin A		
<b>Immunogen:</b> KLH conjugated synthetic peptide derived from rat Cystatin A: 1-80/103.		
<b>Purification:</b> affinity purified by Protein A		
<b>Concentration:</b> 1mg/ml		
<b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		<b>Predicted MW.:</b> 12 kDa
<b>Background:</b> The cystatin superfamily encompasses proteins that contain multiple cystatin-like sequences. Some of the members are active cysteine protease inhibitors, while others have lost or perhaps never acquired this inhibitory activity. There are three inhibitory families in the superfamily, including the type 1 cystatins (stefins), type 2 cystatins, and kininogens. This gene encodes a stefin that functions as a cysteine protease inhibitor, forming tight complexes with papain and the cathepsins B, H, and L. The protein is one of the precursor proteins of cornified cell envelope in keratinocytes and plays a role in epidermal development and maintenance. Stefins have been proposed as prognostic and diagnostic tools for cancer. [provided by RefSeq, Jul 2008]		<b>Subcellular Location:</b> Cytoplasm

**— SELECTED CITATIONS —**

- **[IF=3.06]** Wei, Wei. et al. Tamoxifen Activates Dormant Primordial Follicles in Mouse Ovaries. Reprod Sci. 2022 Feb;;1-9 IHC ;Mouse. 35212933