

bs-12015R**[Primary Antibody]****GPR18 Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) Reactivity: Human (predicted: Mouse, Rat, Rabbit, Pig, Horse) Predicted MW.: 38 kDa Subcellular Location: Cell membrane
Clonality: Polyclonal		
GeneID: 2841	SWISS: Q14330	
Target: GPR18		
Immunogen: KLH conjugated synthetic peptide derived from human G protein coupled receptor 18: 131-230/331. < Extracellular >		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: G protein-coupled receptors (GPRs), also known as seven transmembrane receptors, heptahelical receptors or 7TM receptors, comprise a superfamily of proteins that play a role in many different stimulus-response pathways. G protein coupled receptors translate extracellular signals into intracellular signals (G protein activation) and they respond to a variety of signaling molecules, such as hormones and neurotransmitters. GPR18 is a 331 amino acid multi-pass membrane protein that belongs to the G-protein coupled receptor family. Expressed abundantly in spleen and testis, GPR18 functions as a receptor for N-arachidonyl glycine and is thought to contribute to the regulation of the immune system. GPR18 activity is mediated by G proteins that specifically inhibit adenylyl cyclase.		

— VALIDATION IMAGES —

Paraformaldehyde-fixed, paraffin embedded (Human brain glioma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (GPR18) Polyclonal Antibody, Unconjugated (bs-12015R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

— SELECTED CITATIONS —

- **[IF=5.858]** Matthieu Bardin. et al. The resolvin D2 – GPR18 axis is expressed in human coronary atherosclerosis and transduces atheroprotection in apolipoprotein E deficient mice. BIOCHEM PHARMACOL. 2022 Jul;201:115075 IHC ;Human. 35525326