

bs-12035R**[Primary Antibody]**

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

Paralemmin 1 Rabbit pAb**— DATASHEET —**

Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) ICC/IF (1:100-500) ELISA (1:5000-10000) Reactivity: (predicted: Human, Mouse, Rat, Pig, Sheep, Cow, Dog) Predicted MW.: 42 kDa Subcellular Location: Cell membrane ,Cytoplasm
Clonality: Polyclonal		
GeneID: 5064	SWISS: O75781	
Target: Paralemmin 1		
Immunogen: KLH conjugated synthetic peptide derived from human Paralemmin 1: 130-230/387.		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: Paralemmin, also called Paralemmin-1 or PALM, is a widely expressed peripheral membrane protein that is involved in cell structure and shape. A hydrophobic protein, Paralemmin is anchored to the cytoplasmic side of the cell membrane via di-palmitoylation and prenylation of its C-terminal cysteine cluster. Functioning at the synapse to regulate neuronal plasticity and plasma membrane dynamics, Paralemmin can bind to the dopamine receptor D3, thereby reducing D3 expression and subsequent adenylate cyclase activity. Overexpression of Paralemmin induces fibroblasts to extend long filopodia and to assume extreme cell shapes, suggesting involvement in the formation and stabilization of the plasma membrane. Two isoforms of Paralemmin exists due to alternative splicing events.		