
GCNT2 Rabbit pAb

Catalog Number: bs-13316R

Target Protein: GCNT2

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000)

Reactivity: Human, Mouse, Rat (predicted:Horse)

Predicted MW: 46 kDa

Entrez Gene: 2651

Swiss Prot: Q8N0V5

Source: KLH conjugated synthetic peptide derived from human GCNT2: 151-250/400.


Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.


Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: Belonging to the glycosyltransferase 14 family, GCNT2 (glucosaminyl (N-acetyl) transferase 2, I-branching enzyme (I blood group)), also known as II, N-acetylglucosaminyltransferase, IGNT, CCAT, ULG3, GCNT5, GCNT2C or NACGT1, is a 400 amino acid glycosyltransferase that localizes to the Golgi apparatus. Other members of the glycosyltransferase 14 family include GCNT1, GCNT3, GCNT4, GCNT6 and GCNT7. A single-pass type II membrane protein, GCNT2 functions as a branching enzyme known as beta-1,6-N-acetylglucosaminyltransferase, which converts fetal i antigen to adult I antigen in erythrocytes during embryonic development. With expression levels increasing significantly during oncogenesis and development, GCNT2 is found at highest levels in adult prostate and fetal brain, and is found at low levels in heart, small intestine, colon, brain, pancreas and kidney.

VALIDATION IMAGES



Sample: Placenta (Mouse) Lysate at 40 ug Primary: Anti-GCNT2(bs-10196R) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 46 kD Observed band size: 56 kD



Sample: Lane 1: Rat Cerebrum tissue lysates Lane 2: Human A431 cell lysates Primary: Anti-GCNT2 (bs-13316R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 46 kDa Observed band size: 46 kDa