bs-0599R

[Primary Antibody]

Bioss

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

E2F1 Rabbit pAb

- DATASHEET -

Host: Rabbit **Isotype:** IgG

Clonality: Polyclonal

GeneID: 1869 **SWISS:** Q01094

Target: E2F1

Immunogen: KLH conjugated synthetic peptide derived from human E2F1:

101-180/437.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: E2F's are DNA binding proteins, which associate with negative

regulators, such as the retinoblastoma p107 protein, resulting in an altered rate of gene transcription. The E2F proteins contain several evolutionally conserved domains found in most members of the family. These domains include a DNA binding domain, a dimerization domain which determines interaction with the differentiation regulated transcription factor proteins (DP), a transactivation domain enriched in acidic amino acids, and a tumor suppressor protein association domain which is embedded within the transactivation domain. This protein and another 2 members, E2F2 and E2F3, have an additional cyclin binding domain. E2F1 is proposed to be involved in several cellular processes that range from tumor suppressor, cell progression and oncogenesis. E2F1 overexpression can also drive cells into

apoptosis.

Applications: WB (1:500-2000)

IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) ICC/IF (1:100-500)

Reactivity: Human, Mouse

(predicted: Rat, Rabbit, Pig,


Sheep, Cow, Chicken,

Horse)


Predicted 46 kDa

Subcellular Nucleus


VALIDATION IMAGES




Sample: Lane 1: Hela (Human) Cell Lysate at 30 ug Lane 2: A673 (Human) Cell Lysate at 30 ug Lane 3: Molt-4 (Human) Cell Lysate at 30 ug Lane 4: U251 (Human) Cell Lysate at 30 ug Primary: Anti-E2F1 (bs-0599R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 46 kD Observed band size: 58 kD



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (E2F1) Polyclonal Antibody, Unconjugated (bs-0599R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.



HepG2 cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Antibody incubation with (E2F1) polyclonal Antibody, Unconjugated (bs-0599R) 1:100, 90 minutes at 37°C; followed by a conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei



HepG2 cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Antibody incubation with (E2F1) polyclonal Antibody, Unconjugated (bs-0599R) 1:100, 90 minutes at 37°C; followed by a conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei.

- SELECTED CITATIONS -

- [IF=4.85] Junjie Kong. et al. Silencing of RAB42 down-regulated PD-L1 expression to inhibit the immune escape of hepatocellular carcinoma cells through inhibiting the E2F signaling pathway. CELL SIGNAL. 2023 Apr;:110692 ChIP; Human. 37116555
- [IF=4.26] Sopariwala et al. Long-term PGC1β overexpression leads to apoptosis, autophagy and muscle wasting. (2017) Sci.Re. 7:10237 WB; Mouse. 28860475
- [IF=2.74] Liu, Jianhui, et al. "Silica nanoparticle exposure inducing granulosa cell apoptosis and follicular atresia in female Balb/c mice." Environmental Science and Pollution Research (2017): 1-12. WB; Mouse. 29151191
- [IF=3.332] Muhammad T et al. Aloperine in combination with therapeutic adenoviral vector synergistically suppressed the growth of non-small cell lung cancer. J Cancer Res Clin Oncol. 2020 Feb 22. WB; Human. 32088783
- [IF=2.16] Guo, Fangzi, et al. "Endosulfan induces apoptosis by activating the negative regulation pathway of cell cycle and death receptor pathway in spermatogenic cells." Toxicology Research (2017). WB; Rat. 30090493