

bs-1438R

[Primary Antibody]

Hck Rabbit pAb



www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) ICC/IF (1:100-500) ELISA (1:5000-10000) Reactivity: (predicted: Human, Mouse, Rat, Rabbit, Sheep, Cow, Chicken, Horse) Predicted MW.: 58 kDa Subcellular Location: Cell membrane ,Cytoplasm ,Nucleus
Clonality: Polyclonal		
GeneID: 3055	SWISS: P08631	
Target: Hck		
Immunogen: KLH conjugated synthetic peptide derived from human Hck: 301-450/526.		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: HCK is a tyrosine kinase that is predominantly expressed in hemopoietic cell types. It may help couple the Fc receptor to the activation of the respiratory burst. In addition, it may play a role in neutrophil migration and in the degranulation of neutrophils.		

— SELECTED CITATIONS —

- **[IF=2.245]** Pengwei Wang. et al. Deoxynivalenol Induces Inflammation in the Small Intestine of Weaned Rabbits by Activating Mitogen-Activated Protein Kinase Signaling. Front Vet Sci. 2021; 8: 632599 WB,IHC ;Rabbit. 33604367