

bs-11118R**[Primary Antibody]****PCDHA4 Rabbit pAb**

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) ICC/IF (1:100-500) ELISA (1:5000-10000) Reactivity: (predicted: Human, Mouse, Rat, Sheep, Cow, Dog, Horse) Predicted MW.: 99 kDa Subcellular Location: Cell membrane
Clonality: Polyclonal		
GeneID: 56144	SWISS: Q9UN74	
Target: PCDHA4		
Immunogen: KLH conjugated synthetic peptide derived from human PCDHA4: 301-400/947. < Extracellular >		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: Members of the cadherin-related neuronal receptor (CNR) family, including PCDHA4 (also known as CNR1) and PCDHA6 (also known as CNR2), comprise a novel subfamily within the cadherin superfamily of adhesion molecules. The cadherin-related neuronal receptor proteins form a complex with Fyn, a protein tyrosine kinase that is involved in building brain networks and determining patterns of behavior. Cadherin-related neuronal receptor 1 and 2 were discovered during a search for receptor molecules to the Fyn signaling pathway in the mammalian brain. Members of the cadherin superfamily are Ca ²⁺ -dependent adhesion molecules that function to mediate cell-cell binding critical to the maintenance of tissue structure and morphogenesis. The PCDHA4 and PCDHA6 extracellular domains contain six cadherin repeats that mediate Ca ⁺⁺ -dependent cell adhesion, while the cytoplasmic domains are not homologous with other cadherins.		