

---

## phospho-MST4 + MST3 + STK25 (Thr178 + Thr190 + Thr174) Rabbit pAb

Catalog Number: bs-10441R

Target Protein: phospho-MST4 + MST3 + STK25 (Thr178 + Thr190 + Thr174)

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000), IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500)

Reactivity: Human, Mouse, Rat (predicted: Sheep, Cow, Horse)

Predicted MW: 48 kDa

Entrez Gene: 10494

Swiss Prot: O00506

Source: KLH conjugated synthesised phosphopeptide derived from human MST4/MST3/STK25 around the phosphorylation site of Thr178/Thr190/Thr174: RN(p-T)FV.

Purification: affinity purified by Protein A


Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.


**Background:** Novel human Ste20-related kinase Mst4 is biologically active in the activation of MEK/ERK pathway and in mediating cell growth and transformation. It is pro apoptotic and is highly expressed in placenta, thymus, and peripheral blood leukocytes. Interaction with Golgi matrix protein GOLGA2 results in autophosphorylation on Thr-178, possibly as a consequence of stabilization of dimer formation. This may also be activated by C terminal cleavage. MST3 or Mammalian Sterile 20-like kinase 3 is a member of the germinal center kinase-III family. MST3 contains a conserved kinase domain at its NH(2)-terminus and a regulatory domain at its COOH-terminus. Caspase-mediated cleavage of the regulatory domain of MST3 activates its intrinsic kinase activity and leads to nuclear translocation. Expression of COOH-terminal truncated MST3 in cells results in DNA fragmentation and induction of apoptosis. It can inhibit cell migration in a fashion dependent on autophosphorylation and can regulate paxillin phosphorylation through tyrosine phosphatase PTP-PEST. Mitogen activated protein kinase cascades have been conserved throughout evolution. In mammals, these cascades allow responses to complex stimuli such as growth factors and inflammatory cytokines. In yeast, STK25 functions upstream of the

## MAPK cascade.


### VALIDATION IMAGES




Sample: U-87MG(Human) Cell Lysate at 30 ug Primary: Anti-phospho-MST4 + MST3 + STK25 (Thr178 + Thr190 + Thr174) (bs-10441R) at 1/500 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution  
Predicted band size: 48 kD Observed band size: 39 kD




Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-phospho-MST4+MST3+STK25 (Thr178+Thr190+Thr174) Polyclonal Antibody, Unconjugated(bs-10441R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining




Paraformaldehyde-fixed, paraffin embedded (mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (phospho-MST4 + MST3 + STK25 (Thr178 + Thr190 + Thr174) ) Polyclonal Antibody, Unconjugated (bs-10441R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Human kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (phospho-MST4 + MST3 + STK25 (Thr178 + Thr190 + Thr174) ) Polyclonal Antibody, Unconjugated (bs-10441R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (phospho-MST4 + MST3 + STK25 (Thr178 + Thr190 + Thr174) ) Polyclonal Antibody, Unconjugated (bs-10441R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Human stomach); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (phospho-MST4 + MST3 + STK25 (Thr178 + Thr190 + Thr174) ) Polyclonal Antibody, Unconjugated (bs-10441R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.