bs-13979R

[Primary Antibody]

COPG2 Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

- DATASHEET		400-901-9800
Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000)
Clonality: Polyclonal		IHC-P (1:100-500) IHC-F (1:100-500)
GenelD: 26958	SWISS: Q9UBF2	IF (1:100-500)
Target: COPG2		ICC/IF (1:100-500)
Immunogen: KLH conj 1-100/87 Purification: affinity p	jugated synthetic peptide derived from human COPG2: 1. urified by Protein A	Reactivity: (predicted: Human, Mouse, Rat, Cow)
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Predicted _{98 kDa} MW.: ^{98 kDa} Subcellular _{Outoplasm}
Background: The coatomer is a cytosolic protein complex that binds to dilysine motifs and reversibly associates with Golgi non-clathrin-coated vesicles, which further mediate biosynthetic protein transport from the ER, via the Golgi up to the trans Golgi network. Coatomer complex is required for budding from Golgi membranes, and is essential for the retrograde Golgi-to-ER transport of dilysine- tagged proteins. In mammals, the coatomer can only be recruited by membranes associated to ADP-ribosylation factors (ARFs), which are small GTP-binding proteins; the complex also influences the Golgi structural integrity, as well as the processing, activity, and endocytic recycling of LDL receptors.		rom ed ces