bs-10710R

[Primary Antibody]

Bioss ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn techsupport@bioss.com.cn

400-901-9800

– DATASHEET –

Host: Rabbit Isotype: IgG

Clonality: Polyclonal
Target: I-309/CCL1

Immunogen: KLH conjugated synthetic peptide derived from mouse I-309/CCL1:

31-80/92.

I-309/CCL1 Rabbit pAb

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: This gene is one of several cytokine genes clustered on the q-arm of chromosome 17. Cytokines are a family of secreted proteins

involved in immunoregulatory and inflammatory processes. The protein encoded by this gene is structurally related to the CXC

subfamily of cytokines. Members of this subfamily are

characterized by two cysteines separated by a single amino acid. This cytokine is secreted by activated T cells and displays

chemotactic activity for monocytes but not for neutrophils. It binds to the chemokine receptor CCR8. [provided by RefSeq, Jul 2008].

Applications: IHC-P (1:100-500)

IHC-F (1:100-500) IF (1:100-500) ICC/IF (1:100-500) ELISA (1:5000-10000)

Reactivity: (predicted: Mouse, Rat)

Predicted MW.: 10 kDa

Subcellular Location: Secreted

- SELECTED CITATIONS -

• [IF=3.269] Mengxue Zhang. et al. Identification and validation of potential novel biomarkers for oral squamous cell carcinoma. Bioengineered. 2021;12(1):8845-8862 IF,IHC; Human,Mouse. 34606406