## bs-10100R

## [ Primary Antibody ]

# BIOSS

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

# HSP90 alpha Rabbit pAb

- DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**GenelD:** 3320 **SWISS:** P07900

Target: HSP90 alpha

Immunogen: KLH conjugated synthetic peptide derived from human HSP90

alpha: 401-500/732.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: Hsp90 (heat shock protein 90) is a molecular chaperone and is one

of the most abundant proteins in unstressed cells. It is an ubiquitous molecular chaperone found in eubacteria and all branches of eukarya, but it is apparently absent in archaea. Whereas cytoplasmic Hsp90 is essential for viability under all conditions in eukaryotes, the bacterial homologue HtpG is

dispensable under non-heat stress conditions.

Applications: WB (1:500-2000)

IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) ICC/IF (1:100-500)

Reactivity: Human, Mouse


(predicted: Rat, Rabbit, Pig, Sheep, Cow, Chicken, Dog,

Horse)


Predicted MW.: 81 kDa

**Subcellular Location:** Cell membrane ,Cytoplasm


#### VALIDATION IMAGES




Sample: Lane 1: Mouse Testis tissue lysates Lane 2: Mouse Placenta tissue lysates Lane 3: Human U251 cell lysates Lane 4: Human Molt-4 cell lysates Primary: Anti-HSP90 alpha (bs-10100R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 81 kD Observed band size: 90 kD



Sample: Testis (Mouse) Lysate at 40 ug Primary: Anti-HSP90 alpha (bs-10100R) at 1/500 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 81 kD Observed band size: 98 kD



Paraformaldehyde-fixed, paraffin embedded (mouse testis tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (HSP90 alpha) Polyclonal Antibody, Unconjugated (bs-10100R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.



Hela cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Antibody incubation with (HSP90 alpha) polyclonal Antibody, Unconjugated (bs-10100R) 1:100, 90 minutes at 37°C; followed by a conjugated Goat Anti-Rabbit IgG antibody at

### - SELECTED CITATIONS -

- [IF=6.3] Suqin Zhu. et al. Oleanolic Acid Up-Regulated UGT1A1 and Antagonized Inflammation by Affecting the Binding of PXR and PKCα to HSP90α and SRC1. PHYTOTHER RES. 2025 May;: WB,IF,CoIP; Human. 40420319
- [IF=3.24] Ding, Zhujin, et al. "Expression of heat shock protein 90 genes during early development and infection in< i> Megalobrama amblycephala</i> and evidence for adaptive evolution in teleost." Developmental & Comparative Immunology (2013). WB; 23954722