

**bs-8615R****[ Primary Antibody ]****Bioss**  
**ANTIBODIES**

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

**Ribonuclease 3/ECP Rabbit pAb****— DATASHEET —**


<b>Host:</b> Rabbit <b>Clonality:</b> Polyclonal <b>GeneID:</b> 6037 <b>Target:</b> Ribonuclease 3/ECP <b>Immunogen:</b> KLH conjugated synthetic peptide derived from human Ribonuclease 3: 45-160/160. <b>Purification:</b> affinity purified by Protein A <b>Concentration:</b> 1mg/ml <b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. <b>Background:</b> ECP is a 160 amino acid potent cytotoxic secretory protein that belongs to the pancreatic ribonuclease family. ECP localizes to the granule matrix of the eosinophil and is involved in the immune response system. ECP is a cytotoxin, neurotoxin and helminthotoxin that is secreted by activated human eosinophils. ECP exhibits anti-parasitic, antibacterial and ribonucleolytic activities. It has been suggested that ECP induces the neurotoxic effect known as the Gordon phenomenon, a syndrome manifested by ataxia, muscular rigidity, paralysis, and tremor that may lead to death. ECP is considered a marker of eosinophilic inflammation and high levels have been found in cases of active asthma and other allergic diseases	<b>Isotype:</b> IgG  <b>SWISS:</b> P12724	<b>Applications:</b> <b>WB</b> (1:500-2000) <b>ICC/IF</b> (1:100-500)  <b>Reactivity:</b> Human  <b>Predicted MW.:</b> 16 kDa  <b>Subcellular Location:</b> Cytoplasm
--	---	--

**— VALIDATION IMAGES —**

Western blot analysis of lysate from human spleen tissue lysate, using RNASE3 Antibody (C-term)(Cat. #bs-8615R). AP13557b was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysate at 35ug.



RNASE3 Antibody (C-term) (Cat. #bs-8615R) western blot analysis in NCI-H460 cell line lysates (35ug/lane). This demonstrates the RNASE3 antibody detected the RNASE3 protein (arrow).



Confocal immunofluorescent analysis of RNASE3 Antibody (C-term) (Cat#bs-8615R) with NCI-H460 cell followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (green). DAPI was used to stain the cell nuclear (blue).

**— SELECTED CITATIONS —**

- **[IF=12.05]** Fujisawa, Daisuke, et al. "Expression of Mas-related gene X2 on mast cells is upregulated in the skin of patients with severe chronic urticaria." *Journal of Allergy and Clinical Immunology* (2014). IHC ;="Human". 24954276
- **[IF=5.3]** Seo-Hee Kim. et al. The effect of apigenin, an aryl hydrocarbon receptor antagonist, in Phthalate-Exacerbated eosinophilic asthma model. *JOURNAL OF CELLULAR AND MOLECULAR MEDICINE*. 2023 Jun;27(13):1900-1910 WB ;Mouse. 37315181
- **[IF=4.7]** Francesca Bertolini. et al.Vascular remodeling and TSLP/angiogenin overexpression in severe mixed asthma.*RESPIRATORY RESEARCH*.2025 Feb 28;26(1):78. Western blot ;Human. 40022191
- **[IF=4.082]** Yoshimasa Imoto. et al. Enhanced 15-Lipoxygenase 1 Production is Related to Periostin Expression and

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Eosinophil Recruitment in Eosinophilic Chronic Rhinosinusitis. Biomolecules. 2020 Nov;10(11):1568 IF ;Human. 33218117

- **[IF=2.36]** Inada, Noriko, et al. "Histamine H 1 and H 4 receptor expression on the ocular surface of patients with chronic allergic conjunctival diseases." Allergology International (2017). ICC ;="Human". 28391980