

---

## Oligophrenin 1 Rabbit pAb

Catalog Number: bs-20427R

Target Protein: Oligophrenin 1

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500)

Reactivity: Mouse, Rat (predicted:Human, Sheep, Dog, Horse)

Predicted MW: 92 kDa

Entrez Gene: 4983

Swiss Prot: O60890

Source: KLH conjugated synthetic peptide derived from human Oligophrenin 1: 701-802/802.

Purification: affinity purified by Protein A


Storage: Preservative: 0.02% Proclin300, Constituents: 1% BSA, 0.01M PBS, pH7.4.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.


**Background:** Ras p21 can exist in either a physiologically quiescent GDP-binding state or a GTP-binding signal-emitting state (1,2). Interaction of Ras p21 with GTPase activating protein (GAP) can increase the rate of hydrolysis of Ras p21-bound GTP by as much as 1000-fold (3). In mitogenically activated and tyrosine kinase-transformed cells, Ras GAP forms a complex with a protein designated p190 (4). At its amino terminus, p190 contains sequence motifs characteristic of all known GTPases, whereas the carboxy terminus contains sequences similar to those found in the Bcr gene product, n-chimerin and Rho GAP, all of which exhibit intrinsic GAP activity (4,5). Oligophrenin-1 is an additional protein with GTPase activating activity. Oligophrenin-1 is a RhoGAP protein that stimulates GTP hydrolysis of Rho subfamily members and is involved in cell migration, morphogenesis and axon outgrowth (6).

### VALIDATION IMAGES

---



Paraformaldehyde-fixed, paraffin embedded (rat lung tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (NFKB p65) Polyclonal Antibody, Unconjugated (bs-20355R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (mouse brain tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Oligophrenin 1) Polyclonal Antibody, Unconjugated (bs-20427R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.