## bs-12363R

## [ Primary Antibody ]


## FLRT3 Rabbit pAb



sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

- DATASHEET 400-901-9800		400-901-9800
Host: Rabbit	<b>lsotype:</b> IgG	Applications: WB (1:500-2000)
Clonality: Polyclonal		<b>ELISA</b> (1:5000-10000)
GenelD: 23767	SWISS: Q9NZU0	Reactivity: Rat (predicted: Human)
Target: FLRT3		
Purification: affinity purified by	Protein A	
Concentration: 1mg/ml		Predicted MW.: <sup>70 kDa</sup>
<b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Subcellular Location: <sup>Cell</sup> membrane ,Cytoplasm
that contains one repeats and belon protein (FLRT) fam pancreas, brain, p be involved in rece both cell adhesior encoding mouse F fusion and endode important for prop exists as multiple	no acid single-pass type I membrane prote fibronectin type-III domain and ten leucine gs to the fibronectin leucine rich transmen ily. Expressed in heart, liver, lung, kidney, lacenta and skeletal muscle, FLRT3 is thouge ptor signaling events and may play a role i neurite outgrowth. Defects in the gene LRT3 may lead to ventral closure, headfold erm migration defects, suggesting that FLR per cell differentiation and development. Fl alternatively spliced isoforms that are enco naps to human chromosome 20.	e-rich nbrane ght to in 1 T3 is LRT3

## - VALIDATION IMAGES -



Sample: Lane 1: Rat Cerebrum tissue lysates Lane 2: Rat Lung tissue lysates Primary: Anti-FLRT3 (bs-12363R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 70 kDa Observed band size: 73 kDa