bs-12426R

[Primary Antibody]

Bioss ANTIBODIES

phospho-CSNK1G 1+2+3 (Tyr263) Rabbit pAb

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

- DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GenelD: 53944 SWISS: Q9HCP0

Target: phospho-CSNK1G 1+2+3 (Tyr263)

Immunogen: KLH conjugated synthesised phosphopeptide derived from human

Casein kinase I isoforms gamma 1+2+3 around the

phosphorylation site of Tyr263: ER(p-Y)QK.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: Casein kinases are operationally defined by their preferential utilization of acidic proteins such as caseins as substrates. It can

utilization of acidic proteins such as caseins as substrates. It can phosphorylate a large number of proteins. Participates in Wnt

signaling.

Applications: WB (1:500-2000)

IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500)


Reactivity: Mouse, Rat

(predicted: Human, Pig, Cow, Dog, Horse)

Predicted MW.: 49 kDa

Subcellular Cytoplasm Location:


VALIDATION IMAGES



Sample: Brain (Mouse) Lysate at 40 ug Primary: Anti-phospho-CSNK1G 1+2+3 (Tyr263) (bs-12426R) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 49 kD Observed band size: 49 kD



Sample: Lymph node (Mouse) Lysate at 40 ug Primary: Anti-phospho-CSNK1G 1+2+3 (Tyr263) (bs-12426R) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 49 kD Observed band size: 49 kD



Paraformaldehyde-fixed, paraffin embedded (rat brain tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (CSNK1G 1+2+3 (Tyr263)) Polyclonal Antibody, Unconjugated (bs-12426R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.