

bs-12376R**[Primary Antibody]****FAM20C Rabbit pAb****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500)
Clonality: Polyclonal		IHC-F (1:100-500)
GeneID: 56975	SWISS: Q8IXL6	IF (1:100-500)
Target: FAM20C		Reactivity: Rat (predicted: Human, Mouse, Sheep, Cow, Chicken, Dog, Horse, Monkey)
Immunogen: KLH conjugated synthetic peptide derived from human FAM20C: 351-450/584.		Predicted MW.: 64 kDa
Purification: affinity purified by Protein A		Subcellular Location: Secreted
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: The FAM20 proteins are a family of secreted proteins that regulate differentiation and function of hematopoietic and other tissues. FAM20C, also known as DMP4 (Dentin matrix protein 4), is a 570 amino acid secreted protein that binds calcium and may play a role in dentin mineralization. Defects in the gene encoding FAM20C are the cause of Raine syndrome (Lethal osteosclerotic bone dysplasia), an autosomal recessive osteosclerotic bone dysplasia, that is characterized by generalized osteosclerosis, microencephaly and craniofacial dysplasia. Usually, affected individuals survive only days or weeks. The mutations of the FAM20C gene include four nonsynonymous base changes and four splice-site changes that have a detrimental affect on splicing.		

— VALIDATION IMAGES —

Paraformaldehyde-fixed, paraffin embedded (Rat liver); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (FAM20C) Polyclonal Antibody, Unconjugated (bs-12376R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.