

bs-12377R**[Primary Antibody]****BioSS**
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

OSR1 + OSR2 Rabbit pAb


— DATASHEET —

<p>Host: Rabbit</p> <p>Clonality: Polyclonal</p> <p>GeneID: 130497</p> <p>Target: OSR1 + OSR2</p> <p>Immunogen: KLH conjugated synthetic peptide derived from human OSR2 and OSR1: 181-280/312.</p> <p>Purification: affinity purified by Protein A</p> <p>Concentration: 1mg/ml</p> <p>Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.</p> <p>Background: The OSR1 (odd skipped related 1) gene encodes a zinc-finger transcription factor that is expressed in the germ ring mesoderm and subsequently in the endoderm and intermediate mesoderm, prior to the expression of definitive kidney or blood/vascular markers. Osr1 expression marks a multi-potent population of intermediate mesoderm that undergoes progressive restriction to an Osr1-dependent nephron progenitor compartment within the mammalian kidney. Osr1 expression is regulated by Runx2 and Ikzf1, which are known as master-gene of osteogenesis and hematopoiesis, respectively.</p>	<p>Isotype: IgG</p> <p>SWISS: Q8TAX0</p>	<p>Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500)</p> <p>Reactivity: Human, Mouse, Rat (predicted: Pig, Sheep, Cow, Dog, Horse, Danio rerio)</p> <p>Predicted MW.: 35 kDa</p> <p>Subcellular Location: Nucleus</p>
---	--	--

— VALIDATION IMAGES —



Paraformaldehyde-fixed, paraffin embedded (human testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (OSR1 + OSR2) Polyclonal Antibody, Unconjugated (bs-12377R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Tissue/cell: mouse embryo tissue ; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-OSR2 Polyclonal Antibody, Unconjugated(bs-12377R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining