bs-12050R

[Primary Antibody]

BIOSS

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

5HT2B Receptor Rabbit pAb

- DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GenelD: 3357 **SWISS:** P41595

Target: 5HT2B Receptor

Immunogen: KLH conjugated synthetic peptide derived from human 5HT2B

Receptor/SR-2B: 1-100/481. < Extracellular >

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: Multiple receptor subtypes of serotonin neurotransmitters with

multiple receptor subtypes of serotomin features with multiple physiologic functions have been recognized. The 5-HT-2 receptors mediate many of the central and peripheral physiologic functions of serotonin. Cardiovascular effects include contraction of blood vessels and shape changes in platelets; central nervous system effects include neuronal sensitization to tactile stimuli and mediation of hallucinogenic effects of phenylisopropylamine

hallucinogens.[supplied by OMIM].

Applications: WB (1:500-2000)


IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500)

Reactivity: Human


Predicted MW.: 54 kDa

Subcellular Location: Cell membrane

VALIDATION IMAGES



Sample: Lane 1: Human HepG2 cell lysates Lane 2: Human SH-SY5Y cell lysates Lane 3: Human MCF-7 cell lysates Lane 4: Human HeLa cell lysates Primary: Anti-5HT2B Receptor (bs-12050R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 54 kDa Observed band size: 60 kDa



Paraformaldehyde-fixed, paraffin embedded (human colon carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (5HT2B Receptor) Polyclonal Antibody, Unconjugated (bs-12050R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.