

**bs-12056R****[ Primary Antibody ]****Bioss**  
**ANTIBODIES**

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

**5HT7 Receptor Rabbit pAb****— DATASHEET —**


<b>Host:</b> Rabbit	<b>Isotype:</b> IgG	<b>Applications:</b> WB (1:500-2000) <b>IHC-P</b> (1:100-500) <b>IHC-F</b> (1:100-500) <b>IF</b> (1:100-500)  <b>Reactivity:</b> Human, Mouse, Rat  <b>Predicted MW.:</b> 54 kDa  <b>Subcellular Location:</b> Cell membrane
<b>Clonality:</b> Polyclonal		
<b>GeneID:</b> 3363	<b>SWISS:</b> P34969	
<b>Target:</b> 5HT7 Receptor		
<b>Immunogen:</b> KLH conjugated synthetic peptide derived from human 5HT7 Receptor/SR-7: 51-150/479.		
<b>Purification:</b> affinity purified by Protein A		
<b>Concentration:</b> 1mg/ml		
<b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
<b>Background:</b> The neurotransmitter, serotonin, is thought to play a role in various cognitive and behavioral functions. The serotonin receptor encoded by this gene belongs to the superfamily of G protein-coupled receptors and the gene is a candidate locus for involvement in autistic disorder and other neuropsychiatric disorders. Three splice variants have been identified which encode proteins that differ in the length of their carboxy terminal ends. [provided by RefSeq, Jul 2008]		

**— VALIDATION IMAGES —**


Sample: Lane 1: Mouse Cerebrum tissue lysates  
 Lane 2: Mouse Cerebellum tissue lysates Lane 3:  
 Rat Cerebrum tissue lysates Lane 4: Rat  
 Cerebellum tissue lysates Primary: Anti-5HT7  
 Receptor (bs-12056R) at 1/1000 dilution  
 Secondary: IRDye800CW Goat Anti-Rabbit IgG at  
 1/20000 dilution Predicted band size: 54 kDa  
 Observed band size: 50 kDa



Sample: Hippocampus (Mouse) Lysate at 40 ug  
 Hippocampus(Rat) Lysate at 40 ug Primary: Anti-  
 5HT7 Receptor (bs-12056R) at 1/1000 dilution  
 Secondary: IRDye800CW Goat Anti-Rabbit IgG at  
 1/20000 dilution Predicted band size: 54 kD  
 Observed band size: 54 kD



Paraformaldehyde-fixed, paraffin embedded  
 (mouse cerebellum); Antigen retrieval by boiling  
 in sodium citrate buffer (pH6.0) for 15min; Block  
 endogenous peroxidase by 3% hydrogen  
 peroxide for 20 minutes; Blocking buffer (normal  
 goat serum) at 37°C for 30min; Incubation with  
 (5HT7 Receptor) Polyclonal Antibody,  
 Unconjugated (bs-12056R) at 1:200 overnight at  
 4°C, followed by operating according to SP  
 Kit(Rabbit) (sp-0023) instructions and DAB  
 staining.



Paraformaldehyde-fixed, paraffin embedded  
 (mouse cerebellum); Antigen retrieval by boiling  
 in sodium citrate buffer (pH6.0) for 15min; Block  
 endogenous peroxidase by 3% hydrogen  
 peroxide for 20 minutes; Blocking buffer (normal  
 goat serum) at 37°C for 30min; Incubation with

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

(5HT7 Receptor) Polyclonal Antibody,  
Unconjugated (bs-12056R) at 1:200 overnight at  
4°C, followed by operating according to SP  
Kit(Rabbit) (sp-0023) instructions and DAB  
staining.

## — SELECTED CITATIONS —

---

- **[IF=1.7]** Seyma Ozsoy. et al. Ondansetron and AS19 attenuate morphine tolerance by modulating serotonin 5-HT3 and 5-HT7 receptor expressions in rat dorsal root ganglia. J RADIAT RES APPL SC. 2023 Dec;16:100682 IHC ;Rat. 10.1016/j.jrras.2023.100682