## [ Primary Antibody ]

## phospho-GARB1 (Ser434) Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

		400-90	JI-9800
Host: Rabbit	<b>lsotype:</b> IgG	Applications:	<b>WB</b> (1:500-2000)
Clonality: Polyclonal			<b>IHC-P</b> (1:100-500)
GenelD: 2560	SWISS: P18505		IF (1:100-500)
Target: GARB1 (Ser434)			ICC/IF (1:100-500)
Immunogen: KLH conjugated synthesised phosphopeptide derived from human GARB1 around the phosphorylation site of Ser434: RA(p-S)QL.		Reactivity: Mouse (predicted: Human,	
Purification: affinity purified by Protein A			Rat, Rabbit, Pig, Sheep,
Concentration: 1mg/ml			cow, norse)
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Predicted 51 kDa MW.: <sup>51 kDa</sup> Subcellular <sub>Cell membrane</sub>	
Background: GAD-65 and GAD-67, glutamate decarboxylases, function to catalyze the production of GABA (g-aminobutyric acid). In the central nervous system GABA functions as the main inhibitory transmitter by increasing a Cl-conductance that inhibits neuronal firing. GABA has been shown to activate both ionotropic (GABAA) and metabotropic (GABAB) receptors as well as a third class of receptors called GABAC. Both GABAA and GABAC are ligand-gated ion channels, however, they are structurally and functionally distinct. Members of the GABAA receptor family include GABAA R alpha 1-6, GABAA R beta 1-3, GABAA R©1-3, GABAA R∂, GABAA R gamma, GABAA R delta 1 and GABAA R delta 2. The GABAB family is composed of GABAB R1 alpha and GABAB R1 beta. GABA T-2 and GABA T-3 (also designated GAT-1, -2 and -3). The GABA transporters function to terminate GABA action.		Location:	