

bs-12064R**[Primary Antibody]****phospho-GARB1 (Ser434) Rabbit pAb**

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) ICC/IF (1:100-500) ELISA (1:5000-10000) Reactivity: Mouse (predicted: Human, Rat, Rabbit, Pig, Sheep, Cow, Horse) Predicted MW.: 51 kDa Subcellular Location: Cell membrane
Clonality: Polyclonal		
GeneID: 2560	SWISS: P18505	
Target: GARB1 (Ser434)		
Immunogen: KLH conjugated synthesised phosphopeptide derived from human GARB1 around the phosphorylation site of Ser434: RA(p-S)QL.		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: GAD-65 and GAD-67, glutamate decarboxylases, function to catalyze the production of GABA (g-aminobutyric acid). In the central nervous system GABA functions as the main inhibitory transmitter by increasing a Cl-conductance that inhibits neuronal firing. GABA has been shown to activate both ionotropic (GABAA) and metabotropic (GABAB) receptors as well as a third class of receptors called GABAC. Both GABAA and GABAC are ligand-gated ion channels, however, they are structurally and functionally distinct. Members of the GABAA receptor family include GABAA R alpha 1-6, GABAA R beta 1-3, GABAA R δ 1-3, GABAA R ρ , GABAA R gamma, GABAA R delta 1 and GABAA R delta 2. The GABAB family is composed of GABAB R1 alpha and GABAB R1 beta. GABA transporters have also been identified and include GABA T-1, GABA T-2 and GABA T-3 (also designated GAT-1, -2 and -3). The GABA transporters function to terminate GABA action.		