

**bs-6376R****[ Primary Antibody ]****DDIT4L Rabbit pAb****Bioss**  
**ANTIBODIES**

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

**— DATASHEET —**

<b>Host:</b> Rabbit	<b>Isotype:</b> IgG	<b>Applications:</b> IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500)  <b>Reactivity:</b> Human, Mouse (predicted: Rat, Rabbit, Pig, Cow, Dog, Horse)  <b>Predicted MW.:</b> 22 kDa  <b>Subcellular Location:</b> Cytoplasm
<b>Clonality:</b> Polyclonal		
<b>GeneID:</b> 115265	<b>SWISS:</b> Q96D03	
<b>Target:</b> DDIT4L		
<b>Immunogen:</b> KLH conjugated synthetic peptide derived from human DDIT4L: 121-193/193.		
<b>Purification:</b> affinity purified by Protein A		
<b>Concentration:</b> 1mg/ml		
<b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
<b>Background:</b> DDIT4L inhibits cell growth by regulating the FRAP1 pathway upstream of the TSC1-TSC2 complex and downstream of AKT1. It is up regulated in atherosclerotic plaques relative to healthy segments of the same artery.		

**— VALIDATION IMAGES —**

Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (DDIT4L) Polyclonal Antibody, Unconjugated (bs-6376R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.