
Phospho-PIM1 (Ser99) Rabbit pAb

Catalog Number: bs-20020R

Target Protein: Phospho-PIM1 (Ser99)

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500)

Reactivity: Mouse (predicted:Human, Rat, Pig, Sheep, Cow, Chicken, Dog, GuineaPig, Horse)

Predicted MW: 44 kDa

Entrez Gene: 5292

Swiss Prot: P11309

Source: KLH conjugated synthesised phosphopeptide derived from human PIM1 around the phosphorylation site of Ser99: IN(p-S)LA.


Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: PIM1 is a member of the serine/threonine kinase PIM oncogene family. PIM1 has been implicated in lymphomagenesis, cell proliferation, apoptosis, differentiation and tumourigenesis. The PIM1 protein kinase is upregulated in prostate cancer. The Pim family serine/threonine protein kinases were first identified in studies examining genes targeted for proviral insertion in murine leukemia virus-induced T lymphomas. Increased levels of Pim kinases predispose cells to lymphomagenesis and enhance the activity of mitogenic proteins such as p100, c-Myb, and Cdc25A. In addition, Pim kinases are also involved in modulation of synaptic strength in neurons and anti-apoptotic signaling in hematopoietic progenitor cells. Pim-3, a member of the proto-oncogene Pim family that expresses serine/threonine kinase activity, shares significant homology with Pim-1 serine/threonine protein kinases. Pim-3 may function as a mediator of synaptic plasticity in the brain and is presumably involved in the anti-apoptosis process and cell cycle progression as well as the proliferation of human hepatoma cell lines. The Pim-3 protein is widely expressed, however no expression is observed in the colon, thymus, or small intestine.

VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (mouse kidney tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (PIM1 (Ser99)) Polyclonal Antibody, Unconjugated (bs-20020R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.