

BH0156**[Primary Antibody]****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

MUC6(ready to use) Mouse mAb**— DATASHEET —**

Host: Mouse	Isotype: IgG2a	Applications: IHC-P IHC-F IF ICC/IF Reactivity: Human Predicted MW.: 298 kDa Subcellular Location: Secreted
Clonality: Monoclonal	CloneNo.: 6H8	
GeneID: 4588	SWISS: Q6W4X9	
Target: MUC6(ready to use)		
Purification: affinity purified by Protein G		
Storage: 0.01M PBS (pH7.4) with 1% BSA and 0.02% Proclin300. Store at 2-8°C for one year. Avoid repeated freeze/thaw cycles.		
Background: This gene encodes a member of the mucin protein family. Mucins are high molecular weight glycoproteins produced by many epithelial tissues. The protein encoded by this gene is secreted and forms an insoluble mucous barrier that protects the gut lumen. [provided by RefSeq, Dec 2016]		

— VALIDATION IMAGES —

Paraformaldehyde-fixed, paraffin embedded (human gastric); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (MUC6) Monoclonal Antibody, Unconjugated (BH0156) overnight at 4°C, followed by operating according to SP Kit(Mouse)(sp-0024) instructions and DAB staining.



araformaldehyde-fixed, paraffin embedded (human gastric carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (MUC6) Monoclonal Antibody, Unconjugated (BH0156) overnight at 4°C, followed by operating according to SP Kit(Mouse)(sp-0024) instructions and DAB staining.