

bs-11131R**[Primary Antibody]****alpha Lactalbumin Rabbit pAb****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000)
Clonality: Polyclonal		Reactivity: Human
GeneID: 3906	SWISS: P00709	
Target: alpha Lactalbumin		
Immunogen: KLH conjugated synthetic peptide derived from human LALBA/alpha Lactalbumin: 61-142/142.		Predicted MW.: 14 kDa
Purification: affinity purified by Protein A		Subcellular Location: Secreted
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: Alpha-lactalbumin is the B protein of lactose synthetase secreted by the mammary epithelial cells. It is a potent Ca ²⁺ -elevating and apoptosis-inducing agent with broad, yet selective, cytotoxic activity. Multimeric ?lactalbumin has been shown to kill all transformed, embryonic and lymphoid cells tested, but not mature epithelial elements. This suggests that milk contributes to mucosal immunity not only by furnishing antimicrobial molecules but also by policing the function of lymphocytes and epithelium. ?lactalbumin may be helpful in discovering the site of origin of metastatic breast tumors. Human lactalbumin contains 123 amino acid residues. Comparison of the 5' flanking sequences of the two Alpha-lactalbumin genes with those of five casein genes reveals the presence of a highly conserved region extending from position -140 to -110 in all seven sequences examined, suggesting a possible regulatory role in the hormonal control or tissue-specific expression of milk protein genes in the mammary gland.		

— VALIDATION IMAGES —

Sample: Lane1:Human milk Lysates
Lane2:Human milk Lysates Primary: Anti- alpha Lactalbumin (bs-11131R) at 1/1000 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 14 kD
Observed band size: 14 kD

— SELECTED CITATIONS —

- **[IF=2.6]** Ning Yu. et al. New insight into the effects of different glycation treatments on the structure and IgG-binding capacity of α -lactalbumin. INT J FOOD SCI TECH. 2024 Jul;; Other ;. 10.1111/ijfs.17344