

bs-6616R**[Primary Antibody]****GATA2 Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —**Host:** Rabbit**Isotype:** IgG**Clonality:** Polyclonal**GeneID:** 2624**SWISS:** P23769**Target:** GATA2**Immunogen:** KLH conjugated synthetic peptide derived from human GATA-2: 351-450/480.**Purification:** affinity purified by Protein A**Concentration:** 1mg/ml**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.


Background: Members of the GATA family share a conserved zinc finger DNA-binding domain and are capable of binding the WGATAR consensus sequence. GATA-1 is erythroid-specific and is responsible for the regulated transcription of erythroid genes. It is an essential component in the generation of the erythroid lineage. GATA-2 is expressed in embryonic brain and liver, HeLa and endothelial cells, as well as in erythroid cells. Studies with a modified GATA consensus sequence, AGATCTTA, have shown that GATA-2 and GATA-3 recognize this mutated consensus while GATA-1 has poor recognition of this sequence. This indicates broader regulatory capabilities of GATA-2 and GATA-3 than GATA-1. GATA-3 is highly expressed in T lymphocytes. GATA-4, GATA-5 and GATA-6 comprise a subfamily of transcription factors. Both GATA-4 and GATA-6 are found in heart, pancreas and ovary; lung and liver tissues exhibit GATA-6, but not GATA-4 expression. GATA-5 expression has been observed in differentiated heart and gut tissues and is present throughout the course of development in the heart. Although expression patterns of the various GATA transcription factors may overlap, it is not yet apparent how the GATA factors are able to discriminate in binding their appropriate target sites.

Applications: WB (1:500-2000)**IHC-P** (1:100-500)**IHC-F** (1:100-500)**IF** (1:100-500)


Reactivity: Mouse, Rat
(predicted: Human, Rabbit, Pig, Sheep, Cow, Chicken, Horse)

Predicted MW.: 53 kDa


Subcellular Location: Nucleus

— VALIDATION IMAGES —

Sample: Lane 1: Mouse Heart tissue lysates
Lane 2: Mouse Liver tissue lysates
Primary: Anti-GATA2 (bs-6616R) at 1/1000 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 53 kDa
Observed band size: 55 kDa



Paraformaldehyde-fixed, paraffin embedded (rat duodenum); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (GATA2) Polyclonal Antibody, Unconjugated (bs-6616R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Mouse placenta); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (GATA2) Polyclonal Antibody, Unconjugated (bs-6616R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

— SELECTED CITATIONS —

- **[IF=3.87]** Ikari, Akira, et al. "Hyperosmolarity - Induced Down - Regulation of Claudin - 2 Mediated by Decrease in PKC β -Dependent GATA - 2 in MDCK Cells." *Journal of Cellular Physiology* (2015). WB ;="Dog". 25825272
- **[IF=3.15]** Yang Zhai. et al. Construction of the optimization prognostic model based on differentially expressed immune genes of lung adenocarcinoma. *Bmc Cancer*. 2021 Dec;21(1):1-13 IHC ;Human. 33648465