

bsm-33094M**[Primary Antibody]**

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

Histone H3 (tri methyl K36) Mouse mAb**— DATASHEET —**

| | | |
|---|---|---|
| Host: Mouse Clonality: Monoclonal GeneID: 8350 | Isotype: IgG1 CloneNo.: 2A12 SWISS: P68431 | Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) |
| Target: Histone H3 (tri methyl K36) | | |
| Immunogen: KLH conjugated synthesised methylpeptide derived from human Histone H3 around the methylation site of Tri Methyl K36: GV(Tri Methyl-K)KP. | | |
| Purification: affinity purified by Protein G | | Reactivity: Human, Mouse, Rat |
| Concentration: 1mg/ml | | |
| Storage: Size : 50ul/100ul/200ul 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Size : 200ug (PBS only) 0.01M PBS Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. | | Predicted MW.: 15 kDa Subcellular Location: Nucleus |
| Background: Modulation of the chromatin structure plays an important role in the regulation of transcription in eukaryotes. The nucleosome, made up of four core histone proteins (H2A, H2B, H3 and H4), is the primary building block of chromatin. The N-terminal tail of core histones undergoes different posttranslational modifications including acetylation, phosphorylation and methylation. These modifications occur in response to cell signal stimuli and have a direct effect on gene expression. In most species, the histone H2B is primarily acetylated at lysines 5, 12, 15 and 20. Histone H3 is primarily acetylated at lysines 9, 14, 18 and 23. Acetylation at lysine 9 appears to have a dominant role in histone deposition and chromatin assembly in some organisms. Phosphorylation at Ser10 of histone H3 is tightly correlated with chromosome condensation during both mitosis and meiosis. | | |

— VALIDATION IMAGES —

Paraformaldehyde-fixed, paraffin embedded (rat liver); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Histone H3 (tri methyl K36)) Monoclonal Antibody, Unconjugated (bsm-33094M) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Mouse)(sp-0024) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (rat colon); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Histone H3 (tri methyl K36)) Monoclonal Antibody, Unconjugated (bsm-33094M) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Mouse)(sp-0024) instructions and DAB staining.