

bs-12395R**[Primary Antibody]****KAT3A / CBP Rabbit pAb**

Bioss
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

DATASHEET

Host: Rabbit
Clonality: Polyclonal
GeneID: 1387
Target: KAT3A / CBP
Immunogen: KLH conjugated synthetic peptide derived from human KAT3A/CBP: 301-400/2442.
Purification: affinity purified by Protein A
Concentration: 1mg/ml
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
 Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Isotype: IgG**SWISS:** Q92793


Applications: IHC-P (1:100-500)
 IHC-F (1:100-500)
 IF (1:100-500)
 Flow-Cyt (1ug/Test)

Reactivity: Human, Mouse
 (predicted: Rat, Pig, Sheep,
 Cow, Chicken, Dog, Horse)


Predicted
MW.: 265 kDa

Subcellular
Location: Cytoplasm ,Nucleus


Background: This gene is ubiquitously expressed and is involved in the transcriptional coactivation of many different transcription factors. First isolated as a nuclear protein that binds to cAMP-response element binding protein (CREB), this gene is now known to play critical roles in embryonic development, growth control, and homeostasis by coupling chromatin remodeling to transcription factor recognition. The protein encoded by this gene has intrinsic histone acetyltransferase activity and also acts as a scaffold to stabilize additional protein interactions with the transcription complex. This protein acetylates both histone and non-histone proteins. This protein shares regions of very high sequence similarity with protein p300 in its bromodomain, cysteine-histidine-rich regions, and histone acetyltransferase domain. Mutations in this gene cause Rubinstein-Taybi syndrome (RTS). Chromosomal translocations involving this gene have been associated with acute myeloid leukemia. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Feb 2009].

VALIDATION IMAGES

Paraformaldehyde-fixed, paraffin embedded (mouse intestine); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (KAT3A CBP) Polyclonal Antibody, Unconjugated (bs-12395R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (human colon carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (KAT3A CBP) Polyclonal Antibody, Unconjugated (bs-12395R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Blank control:Hela. Primary Antibody (green line): Rabbit Anti-KAT3A/CBP antibody (bs-12395R) Dilution: 1ug/Test; Secondary Antibody : Goat anti-rabbit IgG-FITC Dilution: 0.5ug/Test. Protocol The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 90% ice-cold methanol for 20 min at -20°C. The cells were then incubated in 5% BSA to block non-specific protein-protein interactions for 30 min at room temperature. Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.