

---

## KIFC1 Rabbit pAb

Catalog Number: bs-7805R

Target Protein: KIFC1

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000), IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500)

Reactivity: Human, Rat (predicted: Mouse, Pig, Cow, Horse)

Predicted MW: 74 kDa

Entrez Gene: 3833

Swiss Prot: Q9BW19

Source: KLH conjugated synthetic peptide derived from human KIFC1: 401-500/673.

Purification: affinity purified by Protein A


Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.


**Background:** The kinesins constitute a large family of microtubule-dependent motor proteins, which are responsible for the distribution of numerous organelles, vesicles and macromolecular complexes throughout the cell. Individual kinesin members play crucial roles in cell division, intracellular transport and membrane trafficking events including endocytosis and transcytosis. KIFC1 is a 673 amino acid protein that belongs to the kinesin-like family of proteins. KIFC1 localizes to the nucleus and contains a C-terminal kinesin-motor domain. Functioning as a minus-end directed microtubule-dependent motor, KIFC1 works together with NuMA and cytoplasmic Dynein to organize microtubule minus ends at spindle poles. HeLa cells deficient in KIFC1 exhibit multipolar mitotic spindles, suggesting that KIFC1 is essential for bipolar spindle formation.

### VALIDATION IMAGES


---




Sample: Raji Cell (Human) Lysate at 40 ug Primary: Anti-KIFC1 (bs-7805R) at 1/300 dilution Secondary: HRP conjugated Goat-Anti-rabbit IgG (bs-0295G-HRP) at 1/5000 dilution Predicted band size: 74 kD Observed band size: 74 kD



Sample: A549 Cell (Human) Lysate at 40 ug Primary: Anti-KIFC1 (bs-7805R) at 1/300 dilution Secondary: HRP conjugated Goat-Anti-rabbit IgG (bs-0295G-HRP) at 1/5000 dilution Predicted band size: 74 kD Observed band size: 74 kD



Paraformaldehyde-fixed, paraffin embedded (Human pancreatic cancer); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (KIFC1) Polyclonal Antibody, Unconjugated (bs-7805R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (rat kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (KIFC1) Polyclonal Antibody, Unconjugated (bs-7805R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

## PRODUCT SPECIFIC PUBLICATIONS

[IF=3.905] Ya-Lan Wei. et al. Kinesin-14 KIFC1 modulates spindle assembly and chromosome segregation in mouse spermatocytes. Exp

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

