

---

## NFKB1 Rabbit pAb

Catalog Number: bs-1194R

Target Protein: NFKB1

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000), IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500)

Reactivity: Human, Mouse, Rat (predicted:Pig, Cow, Chicken, Dog)

Predicted MW: 48/105 kDa

Entrez Gene: 4790

Swiss Prot: P19838

Source: KLH conjugated synthetic peptide derived from human Nuclear factor NF-kappa-B p50 subunit: 51-100/968.

Purification: affinity purified by Protein A


Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.


**Background:** This gene encodes a 105 kD protein which can undergo cotranslational processing by the 26S proteasome to produce a 50 kD protein. The 105 kD protein is a Rel protein-specific transcription inhibitor and the 50 kD protein is a DNA binding subunit of the NF-kappa-B (NFKB) protein complex. NFKB is a transcription regulator that is activated by various intra- and extra-cellular stimuli such as cytokines, oxidant-free radicals, ultraviolet irradiation, and bacterial or viral products. Activated NFKB translocates into the nucleus and stimulates the expression of genes involved in a wide variety of biological functions. Inappropriate activation of NFKB has been associated with a number of inflammatory diseases while persistent inhibition of NFKB leads to inappropriate immune cell development or delayed cell growth. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Sep 2009].

### VALIDATION IMAGES


---




Sample: Brain(Mouse)lysate at 30ug; Primary: Anti-NFkB p105/p50 (bs-1194R) at 1:300 dilution; Secondary: HRP conjugated Goat-Anti-rabbit IgG(bs-0295G-HRP) at 1: 5000 dilution; Predicted band size: 48/105 kD  
Observed band size: 50 kD




Sample: Lung(Mouse) lysate at 30ug; Primary: Anti-NFkB p105/p50 (bs-1194R) at 1:300 dilution; Secondary: HRP conjugated Goat-Anti-rabbit IgG(bs-0295G-HRP) at 1: 5000 dilution; Predicted band size: 48/105 kD  
Observed band size: 110 kD




Sample: Lane 1: Human Raji cell lysates Lane 2: Human MCF-7 cell lysates Lane 3: Human Jurkat cell lysates  
Primary: Anti-NFkB1 (bs-1194R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 48/105 kDa Observed band size: 135 kDa



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (NFkB p50) Polyclonal Antibody, Unconjugated (bs-1194R) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (NFkB p105) Polyclonal Antibody, Unconjugated (bs-1194R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (rat bladder); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (NFkB p105) Polyclonal Antibody, Unconjugated (bs-1194R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

## PRODUCT SPECIFIC PUBLICATIONS

---

- [IF=17.521] Yi Yan. et al. Nanomedicines Reprogram Synovial Macrophages by Scavenging Nitric Oxide and Silencing CA9 in Progressive Osteoarthritis. *Advanced Science*. 2023 Feb;;2207490 WB ; Mouse . 36748885
- [IF=7.666] Defang Zhou. et al. Musashi-1 and miR-147 Precursor Interaction Mediates Synergistic Oncogenicity Induced by Co-Infection of Two Avian Retroviruses. *CELLS-BASEL*. 2022 Jan;11(20):3312 WB ; Chicken . 36291177
- [IF=6.706] Najla Hajji. et al. The Role of Globularia alypum Explored Ex Vivo In Vitro on Human Colon Biopsies from Ulcerative Colitis Patients. *NUTRIENTS*. 2023 Jan;15(6):1457 IHC ; Human . 36986188
- [IF=6.1] Mohammed Elsayed. et al. Empagliflozin protects against isoprenaline-induced fibrosis in rat heart through modulation of TGF- $\beta$ /SMAD pathway. *LIFE SCI*. 2024 Jan;337:122354 IHC ; Rat . 38110076
- [IF=5.162] Lei Zhao. et al. Proteomic analysis reveals the molecular mechanism of Hippophae rhamnoides polysaccharide intervention in LPS-induced inflammation of IPEC-J2 cells in piglets. *Int J Biol Macromol*. 2020 Dec;164:3294 WB ; Pig . 32888998