

**bs-5781R**

**[ Primary Antibody ]**

## MIG7 Rabbit pAb



www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

### — DATASHEET —

<p><b>Host:</b> Rabbit</p> <p><b>Clonality:</b> Polyclonal</p> <p><b>Target:</b> MIG7</p> <p><b>Immunogen:</b> KLH conjugated synthetic peptide derived from human MIG7: 51-150/207.</p> <p><b>Purification:</b> affinity purified by Protein A</p> <p><b>Concentration:</b> 1mg/ml</p> <p><b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.</p> <p><b>Background:</b> MIG7 is a cysteine-rich protein found in cell membranes and cytoplasm of carcinoma cells. Tumor tissue and blood from over 200 cancer patients have been shown to express MIG7, regardless of tissue origin. In contrast, it has not been detectable in tissues or blood from normal subjects. Expression of MIG7 seems to be restricted to carcinoma cells and putatively to the early placenta (Cytotrophoblast) and it's expression is important for the invasive capabilities of tumor cells.</p>	<p><b>Isotype:</b> IgG</p> <p><b>Applications:</b> <b>WB</b> (1:500-2000) <b>IHC-P</b> (1:100-500) <b>IHC-F</b> (1:100-500) <b>IF</b> (1:100-500) <b>ELISA</b> (1:5000-10000)</p> <p><b>Reactivity:</b> (predicted: Human)</p> <p><b>Predicted MW.:</b> 22 kDa</p> <p><b>Subcellular Location:</b> Cell membrane</p>
--	--

### — SELECTED CITATIONS —

- **[IF=3.309]** Ren K et al. Migration-inducing gene-7 independently predicts poor prognosis of human osteosarcoma and is associated with vasculogenic mimicry. Exp Cell Res. 2018 Aug 1;369(1):80-89. IHC, WB ; Human. 29750896