

bs-8005R**[Primary Antibody]****phospho-DDX5 (Tyr593) Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —

Host: Rabbit Clonality: Polyclonal GeneID: 1655 Target: DDX5 (Tyr593) Immunogen: KLH conjugated synthesised phosphopeptide derived from human DDX5 around the phosphorylation site of Tyr593: QA(p-Y)AY. Purification: affinity purified by Protein A Concentration: 1mg/ml Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. Background: p68 RNA helicase is a nuclear protein that exhibits RNA-dependent ATPase activity. Phosphorylation by protein kinase C inhibits p68 RNA helicase activity. p68 RNA helicase appears to play a role in organ differentiation during development. Furthermore, p68 RNA helicase is expressed in early neural development and in various mesodermal tissues in a number of different chordate embryos. At the cellular level, the expression levels of p68 RNA helicase increases in serum-induced quiescent cell lines. p68 RNA helicase may function as a coactivator for estrogen receptor alpha. Additionally, p68 RNA helicase associates with transcriptional coactivators CBP and p300. p68 RNA helicase localizes to the nucleus under normal conditions. During late telophase, p68 RNA helicase and fibrillarin colocalize to nascent nucleoli. p68 RNA helicase may function as a heterodimer with p72 RNA helicase.	Isotype: IgG SWISS: P17844 Applications: WB (1:500-2000) IHC-P (1:100-500) Reactivity: Human, Rat (predicted: Mouse, Cow, Horse) Predicted MW.: 69 kDa Subcellular Location: Nucleus
---	---

— VALIDATION IMAGES —

Paraformaldehyde-fixed, paraffin embedded (Human kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (phospho-DDX5 (Tyr593)) Polyclonal Antibody, Unconjugated (bs-8005R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (phospho-DDX5 (Tyr593)) Polyclonal Antibody, Unconjugated (bs-8005R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

— SELECTED CITATIONS —

- **[IF=9.995]** Alberto Gualtieri. et al. The RNA helicase DDX5 cooperates with EHMT2 to sustain alveolar rhabdomyosarcoma growth. CELL REP. 2022 Aug;40:111267 WB ;Human. 36044855