bsm-49051M

[Primary Antibody]

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

CPV VP2 Mouse mAb

DATASHEET -

Host: Mouse Isotype: IgG Clonality: Monoclonal CloneNo.: 4G2

Target: CPV VP2

Immunogen: Recombinant Canine parvovirus Capsid protein VP2: 201-584/584.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: Size: 50ul/100ul/200ul

0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Size: 200ug (PBS only)

0.01M PBS


Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: Capsid protein self-assembles to form an icosahedral capsid with a

T=1 symmetry, about 22 nm in diameter, and consisting of 60 copies of two size variants of the capsid proteins, VP1 and VP2, which differ by the presence of an N-terminal extension in the minor protein VP1. The capsid encapsulates the genomic ssDNA. Capsid proteins are responsible for the attachment to host cell receptor TFRC. This attachment induces virion internalization predominantly through clathrin-endocytosis. Binding to the host receptors also induces capsid rearrangements leading to surface exposure of VP1 N-terminus, specifically its phospholipase A2-like region and nuclear localization signal(s). VP1 N-terminus might serve as a lipolytic enzyme to breach the endosomal membrane during entry into host cell. Intracytoplasmic transport involves microtubules and interaction between capsid proteins and host dynein. Exposure of nuclear localization signal probably allows nuclear import of capsids.

VALIDATION IMAGES



Sample: Lane 1: Recombinant CPV VP2 protein, His(bs-49051P) Primary: Anti-CPV VP2 (bsm-49051M) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Mouse IgG at 1/20000 dilution Predicted band size: 65 kDa Observed band size: 47 kDa

Applications: WB (1:500-2000)

ELISA (1:5000-10000)

Reactivity: CPV

Predicted 65 kDa

MW.:

Subcellular Secreted, Nucleus