bsm-52130R

[Primary Antibody]

phospho-Akt1 (Ser473) Recombinant Rabbit mAbA N T B D Www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn

SS

E S

400-901-9800

- DATASHEET		400-501-5800
Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000)
Clonality: Recombinant	CloneNo.: 12A1	IHC-P (1:50-200)
GenelD: 207	SWISS: P31749	IF (1:50-200)
Target: Akt1 (Ser473)		Reactivity: Human Mouse Rat
Immunogen: KLH conjugated Synthesised phosphopeptide derived from human Akt1 around the phosphorylation site of Ser473: QF(p-S)YS.		
Purification: affinity purifie	d by Protein A	
Concentration: 1mg/ml		Predicted MW.: ^{56 kDa}
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Subcellular Cell membrane ,Cytoplasm Location: ,Nucleus
Background: This gene encodes one of the three members of the human AKT serine-threonine protein kinase family which are often referred to as protein kinase B alpha, beta, and gamma. These highly similar AKT proteins all have an N-terminal pleckstrin homology domain, a serine/threonine-specific kinase domain and a C-terminal regulatory domain. These proteins are phosphorylated by phosphoinositide 3-kinase (PI3K). AKT/PI3K forms a key component of many signalling pathways that involve the binding of membrane-bound ligands such as receptor tyrosine kinases, G-protein coupled receptors, and integrin-linked kinase. These AKT proteins therefore regulate a wide variety of cellular functions including cell proliferation, survival, metabolism, and angiogenesis in both normal and malignant cells. AKT proteins are recruited to the cell membrane by phosphatidylinositol 3,4,5-trisphosphate (PIP3) after phosphorylation of phosphatidylinositol 4,5- bisphosphate (PIP2) by PI3K. Subsequent phosphorylation of both threonine residue 308 and serine residue 473 is required for full activation of the AKT1 protein encoded by this gene. Phosphorylation of additional residues also occurs, for example, in response to insulin growth factor-1 and epidermal growth factor. Protein phosphatases act as negative regulators of AKT proteins by dephosphorylating AKT or PIP3. The PI3K/AKT signalling pathway is crucial for tumor cell survival. Survival factors can suppress apoptosis in a transcription-independent manner by activating AKT1 which then phosphorylates and inactivates components of the apoptotic machinery. AKT proteins also participate in the mammalian target of rapamycin (mTOR) signalling pathway which controls the assembly of the eukaryotic translation initiation factor 4F (eIF4E) complex and this pathway, in addition to responding to extracellular signals from growth factors and cytokines, is disregulated in many cancers. Mutations in this gene are associated with multiple types of cancer and excessive tissue growth including Pro		

- VALIDATION IMAGES -



MCF-7 (H) cells were treated with or without

Calyculin A (100nM) for 30 min, 25 µg total protein per lane of cell lysates (see on figure) probed with Phospho-Akt1 (Ser473) monoclonal antibody, unconjugated (bsm-52130R) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.

- SELECTED CITATIONS -

- [IF=3.575] Li Li. et al. Circ_LPAR3 promotes the progression of oral squamous cell carcinoma (OSCC). Biochem Bioph Res Co. 2021 Dec;: IHC ;MOUSE. 34922206
- [IF=3.8] Yaxi Zhou. et al. Silkworm pupa protein-derived peptides alleviate LPS-induced inflammatory response in RAW264.7 macrophage cells through the NF-κB/MAPK/PI3K-AKT signaling pathway. Journal of Agriculture and Food Research. 2024 Jun;16:101165 WB ;MOUSe. 10.1016/j.jafr.2024.101165
- [IF=1.9] Yaji Li. et al. CircMYBL1 suppressed acquired resistance to osimertinib in non-small-cell lung cancer. CANCER GENET-NY. 2024 Jun;284-285:34 WB ;Human. 38626533