#### bsm-33297M

### [ Primary Antibody ]

# BIOSS ANTIBODIES

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

**IHC-P** (1:200-800)

**IHC-F** (1:100-500)

(predicted: Mouse, Sheep,

**IF** (1:100-500)

Applications: WB (1:500-2000)

Reactivity: Human, Rat

Predicted

MW.:

Subcellular Cytoplasm Location:

Cow)

42 kDa

## Actin, alpha skeletal muscle Mouse mAb

- DATASHEET -

Host: Mouse Isotype: IgG
Clonality: Monoclonal CloneNo.: 3E9
GeneID: 58 SWISS: P68133

**Target:** Actin, alpha skeletal muscle **Purification:** affinity purified by Protein G

Concentration: 1mg/ml

Storage: Size: 50ul/100ul/200ul

0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Size: 200ug (PBS only)

0.01M PBS


Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.


**Background:** The product encoded by this gene belongs to the actin family of

proteins, which are highly conserved proteins that play a role in cell motility, structure and integrity. Alpha, beta and gamma actin isoforms have been identified, with alpha actins being a major constituent of the contractile apparatus, while beta and gamma actins are involved in the regulation of cell motility. This actin is an alpha actin that is found in skeletal muscle. Mutations in this gene cause nemaline myopathy type 3, congenital myopathy with excess of thin myofilaments, congenital myopathy with cores, and congenital myopathy with fiber-type disproportion, diseases that lead to muscle fiber defects. [provided by RefSeq, Jul 2008]

## VALIDATION IMAGES




25 ug total protein per lane of various lysates (see on figure) probed with Actin, alpha skeletal muscle monoclonal antibody, unconjugated (bsm-33297M) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.




Paraformaldehyde-fixed, paraffin embedded (human cervical carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Actin, alpha skeletal muscle) Monoclonal Antibody, Unconjugated (bsm-33297M) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Mouse)(sp-0024) instructionsand DAB staining.



boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Actin, alpha skeletal muscle) Monoclonal Antibody, Unconjugated (bsm-33297M-3E9) at 1:400 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.



Paraformaldehyde-fixed, paraffin embedded Human Colon Cancer; Antigen retrieval by



Paraformaldehyde-fixed, paraffin embedded Rat Skeletal muscle; Antigen retrieval by boiling in

boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with Actin, alpha skeletal muscle Monoclonal Antibody,
Unconjugated (bsm-33297M) at 1:200 overnight at 4°C. Followed by conjugated Goat Anti-Mouse IgG antibody (green, bs-0296G-BF488), DAPI (blue, C02-04002) was used to stain the cell nuclei.

sodium citrate buffer (pH6.0) for 15 min; The section was incubated with Actin, alpha skeletal muscle Monoclonal Antibody, Unconjugated (bsm-33297M) at 1:200 overnight at 4°C. Followed by conjugated Goat Anti-Mouse IgG antibody (green, bs-0296G-BF488), DAPI (blue, C02-04002) was used to stain the cell nuclei.