

bsm-52793R**[Primary Antibody]**

Bioss
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

TFRC Recombinant Rabbit mAb**— DATASHEET —**


Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000) Flow-Cyt (1µg/Test) ICC/IF (1:50-200)
Clonality: Recombinant	CloneNo.: 8G6	
GeneID: 7037	SWISS: P02786	
Target: TFRC		
Immunogen: A synthesized peptide derived from human Transferrin receptor: 2-50/760.		
Purification: affinity purified by Protein A		Reactivity: Human, Rat (predicted: Mouse)
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: This gene encodes a cell surface receptor necessary for cellular iron uptake by the process of receptor-mediated endocytosis. This receptor is required for erythropoiesis and neurologic development. Multiple alternatively spliced variants have been identified. [provided by RefSeq, Sep 2015]		
		Predicted MW.: 95 kDa
		Subcellular Location: Secreted ,Cell membrane

— VALIDATION IMAGES —

25 µg total protein per lane of various lysates (see on figure) probed with TFRC monoclonal antibody, unconjugated (bsm-52793R) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.



4% Paraformaldehyde-fixed HeLa (H) cell; Triton X-100 at r.t. for 20 min; Antibody incubation with (TFRC) monoclonal Antibody, unconjugated (bsm-52793R) 1:100, 90 min at 37°C; followed by conjugated Goat Anti-Rabbit IgG antibody (green, bs-40295G-FITC) at 37°C for 90 min, DAPI (blue, C02-04002) was used to stain the cell nuclei. PBS instead of the primary antibody was used as the blank control.



The Jar (H) cells were fixed with 4% PFA (10 min at r.t.) and then permeabilized with 90% ice-cold methanol for 20 min at -20°C, the cells then were incubated in 5% BSA to block non-specific protein-protein interactions (30 min at r.t.). Primary Antibody (green): Rabbit Anti-TFRC antibody (bsm-52793R): 1 µg/10⁶ cells; Secondary Antibody (white blue): Goat anti-Rabbit IgG-FITC (bs-40295G-FITC): 1 µg/test. Isotype Control (orange): Rabbit IgG (bs-0295P). Blank control (black): PBS. Acquisition of 20,000 events was performed.

— SELECTED CITATIONS —

- **[IF=14.7]** Yanhong Xing. et al. Lysosomes finely control macrophage inflammatory function via regulating the release of lysosomal Fe²⁺ through TRPML1 channel. *NAT COMMUN.* 2025 Jan 24;16(1):985. Western blot ;Mouse. 39856099