

**bs-10202R**

**[ Primary Antibody ]**

## HPV16 E7 Rabbit pAb

**Bioss**  
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn


techsupport@bioss.com.cn

400-901-9800


### — DATASHEET —

<b>Host:</b> Rabbit	<b>Isotype:</b> IgG	<b>Applications:</b> <b>WB</b> (1:500-2000) <b>Flow-Cyt</b> (1ug/Test)
<b>Clonality:</b> Polyclonal		<b>Reactivity:</b> Human (predicted: HPV16)
<b>Target:</b> HPV16 E7		<b>Predicted MW.:</b> 11 kDa
<b>Immunogen:</b> KLH conjugated synthetic peptide derived from HPV16 E7 protein: 11-70/98.		
<b>Purification:</b> affinity purified by Protein A		
<b>Concentration:</b> 1mg/ml		
<b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		


### — VALIDATION IMAGES —



Sample: Lane 1: HPV16 E7 fusion protein (bs-49101P) 0.1ug Lysates Lane 2: HPV16 E7 fusion protein (bs-49101P) 0.05ug Lysates  
Primary: Anti-HPV16 E7 (bs-10202R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 11kDa Observed band size: 50kDa



Sample: hela (Human) Cell Lysate at 30 ug mcf-7 (Human) Cell Lysate at 30 ug mda-mb-231 (Human) Cell Lysate at 30 ug siha (Human) Cell Lysate at 30 ug recombinant HPV16-E7 protein (bs-49101P) 100 ng recombinant HPV16-E7 protein (bs-49101P) 50 ng Primary: Anti-HPV16 E7 (bs-10202R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 11 kD Observed band size: 17-20 kD bs-49101P band size: 45 kD



Blank control (black line) :Hela. Primary Antibody (green line): Rabbit Anti-HPV16 E7 antibody (bs-10202R) Dilution:1ug/Test; Secondary Antibody (white blue line) : Goat anti-rabbit IgG-AF488 Dilution: 0.5ug/Test. Isotype control (orange line) : Normal Rabbit IgG Protocol The cells were fixed with 4% PFA (10min at room temperature)and then permeabilized with 90% ice-cold methanol for 20 min at -20°C, The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

### — SELECTED CITATIONS —

- **[IF=4.452]** Siyang Zhang. et al. Nocardia rubra cell-wall skeleton influences the development of cervical carcinoma by promoting the antitumor effect of macrophages and dendritic cells. 2022 Jan 07 WB,IF,IHC ;Mouse,Human. 34994088
- **[IF=1.3]** Yahang Chen. et al. Human papillomavirus type 16 E7 promotes cell viability and migration in cervical cancer by regulating the miR-23a/HOXC8 axis. J OBSTET GYNAECOL. 2024 Feb 13 WB ;Human. 38348790