

bs-11157R**[Primary Antibody]****PCDHGB4 Rabbit pAb**

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

| | |
|--|---|
| <p>Host: Rabbit</p> <p>Clonality: Polyclonal</p> <p>GeneID: 8641</p> <p>Target: PCDHGB4</p> <p>Immunogen: KLH conjugated synthetic peptide derived from human PCDHGB4: 101-200/923. < Extracellular ></p> <p>Purification: affinity purified by Protein A</p> <p>Concentration: 1mg/ml</p> <p>Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.</p> <p>Background: The cadherins are a family of Ca²⁺-dependent adhesion molecules that function to mediate cell-cell binding critical to the maintenance of structure and morphogenesis. Cadherins each contain a large extracellular domain at the N-terminus, which is characterized by a series of five homologous repeats, the most distal of which is thought to be responsible for binding specificity. Cadherin-20, also known as protocadherin gamma B4, is a member of the protocadherin gamma gene cluster. Expressed primarily in fibroblasts, cadherin-20 plays a critical role in neuronal cell-cell connections in the brain. Through its ability to mediate cell-cell adhesion, cadherin-20 is thought to help the body respond to injury by regulating post-traumatic wound healing.</p> | <p>Isotype: IgG</p> <p>Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) ICC/IF (1:100-500) ELISA (1:5000-10000)</p> <p>Reactivity: (predicted: Human, Mouse, Rat, Pig, Cow)</p> <p>Predicted MW.: 97 kDa</p> <p>Subcellular Location: Cell membrane</p> |
|--|---|