

bs-11164R**[Primary Antibody]**

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

Sidekick 1 Rabbit pAb**— DATASHEET —**

Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) ICC/IF (1:100-500) ELISA (1:5000-10000) Reactivity: (predicted: Human, Mouse, Rat, Pig, Sheep, Horse) Predicted MW.: 242 kDa Subcellular Location: Cell membrane
Clonality: Polyclonal		
GeneID: 221935	SWISS: Q7Z5N4	
Target: Sidekick 1		
Immunogen: KLH conjugated synthetic peptide derived from human Sidekick 1: 1501-1600/2213. < Extracellular >		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: Cell adhesion molecules influence cell growth, differentiation, embryogenesis, immune response and cancer metastasis by networking information from the extracellular matrix to the cell. Sidekick-1 (SDK1) is a 2,213 amino acid single-pass membrane protein that functions as a cell adhesion molecule by guiding axonal terminals to specific synapses in developing neurons. Existing as three alternatively spliced isoforms, Sidekick-1 is expressed in retinal neurons and contains thirteen fibronectin type-III domains and six Ig-like C2-type (immunoglobulin-like) domains. Sidekick-1 expression is upregulated in glomeruli of patients with HIV-associated nephropathy, where it leads to podocyte dysfunction. The gene encoding Sidekick-1 maps to human chromosome 7p22.2 and murine chromosome 5 G2.		