

**bsm-33226M****[ Primary Antibody ]****RuBisCO Mouse mAb**

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

**— DATASHEET —**

<b>Host:</b> Mouse	<b>Isotype:</b> IgG	<b>Applications:</b> WB (1:500-2000)
<b>Clonality:</b> Monoclonal	<b>CloneNo.:</b> 3G5	<b>Reactivity:</b> Arabidopsis Thaliana
<b>Target:</b> RuBisCO		
<b>Purification:</b> affinity purified by Protein G		
<b>Concentration:</b> 1mg/ml		
<b>Storage:</b> Size : 50ul/100ul/200ul 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Size : 200ug (PBS only) 0.01M PBS Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		<b>Predicted MW.:</b> 52 kDa
<b>Background:</b> RuBisCO catalyzes two reactions: the carboxylation of D-ribulose 1,5-bisphosphate, the primary event in carbon dioxide fixation, as well as the oxidative fragmentation of the pentose substrate in the photorespiration process. Both reactions occur simultaneously and in competition at the same active site.		<b>Subcellular Location:</b> Cytoplasm

**— SELECTED CITATIONS —**

- **[IF=8.2]** Wen Wang. et al. Characterization of a novel  $\gamma$ -type carbonic anhydrase, Sjy-CA2, in *Saccharina japonica*: Insights into carbon concentration mechanism in macroalgae. INT J BIOL MACROMOL. 2024 Apr;263:130506 WB ;*Saccharina japonica*. 38423426
- **[IF=4.9]** Qilei Zhang. et al. Adaptation of the Invasive Plant *Sphagneticola trilobata* to Flooding Stress by Hybridization with Native Relatives. INT J MOL SCI. 2024 Jan;25(12):6738 WB ;*Sphagneticola trilobata*. 38928441