

bs-9303R

[Primary Antibody]

STIL Rabbit pAb



www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:50-200) ELISA (1:5000-10000) Reactivity: (predicted: Human, Mouse, Rat, Rabbit, Cow) Predicted MW.: 143 kDa Subcellular Location: Cytoplasm
Clonality: Polyclonal		
GeneID: 6491	SWISS: Q15468	
Target: STIL		
Immunogen: KLH conjugated synthetic peptide derived from human STIL: 951-1050/1287.		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: TAL1 disruption at 1p32, a common rearrangement in the T-cell acute lymphoblastic leukemia, usually results in the formation of a SCL interrupting locus (SIL)-TAL1 fusion product. SIL is an immediate early gene whose expression is associated with cell proliferation. The Sil protein exhibits ubiquitous expression in hematopoietic cell lines and tissues. However, Sil protein levels remain tightly regulated during the cell cycle, achieving peak levels in mitosis and diminishing on transition to G1 phase. Overexpression of Sil in primary adenocarcinomas predicts metastatic spread, especially in lung tumors with increased mitotic activity.		

— SELECTED CITATIONS —

- **[IF=2.8]** Shunsuke Ohtsuka. et al. STIL Overexpression Is Associated with Chromosomal Numerical Abnormalities in Non-Small-Cell Lung Carcinoma Through Centrosome Amplification. CURR ONCOL. 2024 Dec;31(12):7936-7949 IHC ;Human. 39727708