
MMP-2 Rabbit pAb

Catalog Number: bs-20705R

Target Protein: MMP-2

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000), IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500), Flow-Cyt (1µg/Test), ICC/IF (1:100)

Reactivity: Human, Mouse, Rat, Rabbit (predicted:Pig, Sheep, Cow, Horse)

Predicted MW: 72 kDa

Entrez Gene: 4313

Swiss Prot: P08253

Source: KLH conjugated synthetic peptide derived from human MMP-2: 81-150/662.


Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.


Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: Proteins of the matrix metalloproteinase (MMP) family are involved in the breakdown of extracellular matrix in normal physiological processes, such as embryonic development, reproduction, and tissue remodeling, as well as in disease processes, such as arthritis and metastasis. Most MMP's are secreted as inactive proproteins which are activated when cleaved by extracellular proteinases. This gene encodes an enzyme which degrades type IV collagen, the major structural component of basement membranes. The enzyme plays a role in endometrial menstrual breakdown, regulation of vascularization and the inflammatory response. Mutations in this gene have been associated with Winchester syndrome and Nodulosis-Arthropathy-Osteolysis (NAO) syndrome. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq].


VALIDATION IMAGES



Sample: SY5Y (Human) Cell Lysate at 30 ug U2os (Human) Cell Lysate at 30 ug U251 (Human) Cell Lysate at 30 ug Siha (Human) Cell Lysate at 30 ug Primary: Anti- MMP-2 (bs-20705R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 62 kD Observed band size: 60 kD



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (MMP-2) Polyclonal Antibody, Unconjugated (bs-20705R) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.



Blank control:Huvec. Primary Antibody (green line): Rabbit Anti-MMP-2 antibody (bs-20705R) Dilution: 2µg /10⁶ cells; Isotype Control Antibody (orange line): Rabbit IgG . Secondary Antibody : Goat anti-rabbit IgG-AF647 Dilution: 1µg /test. Protocol The cells were fixed with 4% PFA (10min at room temperature)and then permeabilized with 90% ice-cold methanol for 20 min at -20°C. The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

PRODUCT SPECIFIC PUBLICATIONS

[IF=4.375] Qiu Lei. et al. MiR-526b-3p Inhibits the Resistance of Glioma Cells to Adriamycin by Targeting MAPRE1. J Oncol.

2022;2022:2402212 WB ; Human . 10.1155/2022/2402212

[IF=2.6] Fu Yuheng. et al. Study on the mechanism of action of Wu Mei Pill in inhibiting rheumatoid arthritis through TLR4-NF-κB pathway.

J ORTHOP SURG RES. 2024 Dec;19(1):1-12 WB ; Rat . 38218891