

**bs-14173R**

**[ Primary Antibody ]**

## UCRC Rabbit pAb



www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

### — DATASHEET —

<b>Host:</b> Rabbit	<b>Isotype:</b> IgG	<b>Applications:</b> <b>IHC-P</b> (1:100-500) <b>IHC-F</b> (1:100-500) <b>IF</b> (1:100-500) <b>ICC/IF</b> (1:100-500) <b>ELISA</b> (1:5000-10000)  <b>Reactivity:</b> (predicted: Human, Mouse, Rat, Rabbit, Pig, Sheep, Cow, Dog)  <b>Predicted MW.:</b> 7 kDa  <b>Subcellular Location:</b> Cytoplasm
<b>Clonality:</b> Polyclonal		
<b>GeneID:</b> 29796	<b>SWISS:</b> Q9UDW1	
<b>Target:</b> UCRC		
<b>Immunogen:</b> KLH conjugated synthetic peptide derived from human UCRC: 2-63/63.		
<b>Purification:</b> affinity purified by Protein A		
<b>Concentration:</b> 1mg/ml		
<b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
<b>Background:</b> UCRC is a subunit of mitochondrial complex III (ubiquinol-cytochrome c reductase; EC 1.10.2.2), which forms the middle segment of the respiratory chain of the inner mitochondrial membrane (Schagger et al., 1995 [PubMed 8592474]).[supplied by OMIM, Mar 2008]		

### — SELECTED CITATIONS —

- **[IF=4.938]** Du J et al. The regulation of skeletal muscle fiber-type composition by betaine is associated with NFATc1/MyoD.J Mol Med (Berl). 2018 Jun 6. WB ;Mouse. 29876588